

MCP-2 Polyclonal Antibody
Catalog # AP73476**Specification**

MCP-2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P80075
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

MCP-2 Polyclonal Antibody - Additional Information**Gene ID** 6355**Other Names**

CCL8; MCP2; SCYA10; SCYA8; C-C motif chemokine 8; HC14; Monocyte chemoattractant protein 2; Monocyte chemotactic protein 2; MCP-2; Small-inducible cytokine A8

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

MCP-2 Polyclonal Antibody - Protein Information**Name** CCL8**Synonyms** MCP2, SCYA10, SCYA8**Function**

Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin. The processed form MCP-2(6-76) does not show monocyte chemotactic activity, but inhibits the chemotactic effect most predominantly of CCL7, and also of CCL2 and CCL5 and CCL8.

Cellular Location

Secreted.

Tissue Location


Highest expression found in the small intestine and peripheral blood cells. Intermediate levels seen in the heart, placenta, lung, skeletal muscle, thymus, colon, ovary, spinal cord and pancreas. Low levels seen in the brain, liver, spleen and prostate

MCP-2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCP-2 Polyclonal Antibody - Images



Western Blot analysis of L929 cells using MCP-2 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of L929 cells using MCP-2 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

MCP-2 Polyclonal Antibody - Background

Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin. The processed form MCP-2(6-76) does not show monocyte chemotactic activity, but inhibits the chemotactic effect most predominantly of CCL7, and also of CCL2 and CCL5 and CCL8.