

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide

Synthetic peptide

Catalog # BP7046a

Specification

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Product Information

Primary Accession

[O43741](#)

Other Accession

[NP_005390](#)**AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Additional Information**

Gene ID 5565

Other Names

5'-AMP-activated protein kinase subunit beta-2, AMPK subunit beta-2, PRKAB2

Target/Specificity

The synthetic peptide sequence is selected from aa 18~33 of human PRKAB2.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Protein Information

Name PRKAB2

Function

Non-catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Beta non-catalytic subunit acts as a scaffold on which the AMPK complex assembles, via its C-terminus that bridges alpha (PRKAA1 or PRKAA2) and gamma subunits (PRKAG1, PRKAG2 or PRKAG3).

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Images

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - Background

The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit may be a positive regulator of AMPK activity. The myristoylation and phosphorylation of this subunit have been shown to affect the enzyme activity and cellular localization of AMPK. This subunit may also serve as an adaptor molecule mediating the association of the AMPK complex.

AMPK beta2 (PRKAB2) Antibody (N-term) Blocking peptide - References

Blume-Jensen P, et al. Nature 2001. 411: 355. Cantrell D, J. Cell Sci. 2001. 114: 1439. Jhiang S Oncogene 2000. 19: 5590. Manning G, et al. Science 2002. 298: 1912. Moller, D, et al. Am. J. Physiol. 1994. 266: C351-C359. Robertson, S. et al. Trends Genet. 2000. 16: 368. Robinson D, et al. Oncogene 2000. 19: 5548. Van der Ven, P, et al. Hum. Molec. Genet. 1993. 2: 1889. Vanhaesebroeck, B, et al. Biochem. J. 2000. 346: 561. Van Weering D, et al. Recent Results Cancer Res. 1998. 154: 271.