

Biotinylated Anti-Rat GRO/KC Antibody
Catalog # ABG10328

Specification

Biotinylated Anti-Rat GRO/KC Antibody - Product Information

Application	WB, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

Biotinylated Anti-Rat GRO/KC Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure recombinant Rat GRO/KC. Anti-Rat GRO/KC specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect Rat GRO/KC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Rat GRO/KC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Rat GRO/KC by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems's Polyclonal Anti-Rat GRO/KC (62-059P) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat GRO/KC.

Direct

To detect Rat GRO/KC by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat GRO/KC.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Biotinylated Anti-Rat GRO/KC Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Rat GRO/KC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Biotinylated Anti-Rat GRO/KC Antibody - Images