

**Anti-Human WNT-3a Antibody**  
Catalog # ABG10443

**Specification**

---

**Anti-Human WNT-3a Antibody - Product Information**

|             |                   |
|-------------|-------------------|
| Application | <b>WB, IHC, E</b> |
| Reactivity  | <b>Human</b>      |
| Host        | <b>Rabbit</b>     |
| Clonality   | <b>Polyclonal</b> |

**Anti-Human WNT-3a Antibody - Additional Information**

**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hWNT-3a. Anti-Human WNT-3a specific antibody was purified by affinity chromatography employing immobilized hWNT-3a matrix.

**WesternBlot**

To detect hWNT-3a by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hWNT-3a is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect hWNT-3a by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human WNT-3a (60-403BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hWNT-3a.

**Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human breast invasive ductal carcinoma. The recommended concentration is 0.125 µg/ml - 0.25 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human WNT-3a Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Anti-Human WNT-3a Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Human WNT-3a Antibody - Images**