

**Anti-Human sRANK Receptor Antibody**  
Catalog # ABG10520**Specification****Anti-Human sRANK Receptor Antibody - Product Information**

Application	<b>WB, IHC, E</b>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**Anti-Human sRANK Receptor Antibody - Additional Information****Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hsRANK Receptor. Anti-Human sRANK Receptor specific antibody was purified by affinity chromatography employing immobilized hsRANK Receptor matrix.

**WesternBlot**

To detect hsRANK Receptor by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsRANK Receptor is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect hsRANK Receptor by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human sRANK Receptor (60-284BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsRANK Receptor.

**Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human breast invasive ductal carcinoma. The recommended concentration is 1.0 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer is recommended. Optimal concentrations and conditions may vary.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human sRANK Receptor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Anti-Human sRANK Receptor Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Human sRANK Receptor Antibody - Images**