

Anti-Murine VEGF Antibody
Catalog # ABG10577**Specification****Anti-Murine VEGF Antibody - Product Information**

Application	WB, IHC, E
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal

Anti-Murine VEGF Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mVEGF. Anti-Murine VEGF specific antibody was purified by affinity chromatography employing immobilized mVEGF matrix.

WesternBlot

To detect mVEGF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mVEGF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect mVEGF by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Murine VEGF (61-127BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mVEGF.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of normal murine kidney. The recommended concentration is 2.5µg/ml for one hour at room temperature. A secondary fluorophore conjugated antibody was used for thirty minutes at room temperature. High pH heat induced antigen retrieval is recommended. Optimal concentrations and conditions may vary.

Neutralization

To yield one-half maximal inhibition [**ND**] of the biological activity of mVEGF (10.00 ng/ml), a concentration of 0.08-0.12 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Murine VEGF Antibody is for research use only and not for use in diagnostic or therapeutic

procedures.

Anti-Murine VEGF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Murine VEGF Antibody - Images