

**Anti-Human Visfatin Antibody**  
Catalog # ABG10584**Specification**

---

**Anti-Human Visfatin Antibody - Product Information**

Application	<b>WB, IHC, E</b>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**Anti-Human Visfatin Antibody - Additional Information****Preparation**

Produced from sera of rabbits immunized with highly pure recombinant Human Visfatin. Anti-Human Visfatin specific antibody was purified by affinity chromatography employing an immobilized Human Visfatin matrix.

**WesternBlot**

To detect Human Visfatin by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Human Visfatin is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect Human Visfatin by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human Visfatin (60-347BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human Visfatin.

**Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human ovary malignant adenocarcinoma. The recommended concentration is 7.5ng/mL -15ng/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary. Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human Visfatin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Anti-Human Visfatin Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Human Visfatin Antibody - Images**