

**Anti-Bcl-2 Picoband Antibody**  
Catalog # ABO10009**Specification****Anti-Bcl-2 Picoband Antibody - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">P10415</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Bcl-2 detection. Tested with WB, IHC-P, Direct ELISA in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Bcl-2 Picoband Antibody - Additional Information**

Gene ID 596

**Other Names**

Apoptosis regulator Bcl-2, BCL2

**Application Details**

Western blot, 0.1-0.5 µg/ml  
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml  
Direct ELISA, 0.1-0.5 µg/ml

**Subcellular Localization**

Mitochondrion outer membrane.

**Tissue Specificity**

Expressed in a variety of tissues.

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

E. coli-derived human Bcl-2 recombinant protein (Position: Q118-E165).

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

## Anti-Bcl-2 Picoband Antibody - Protein Information

**Name** BCL2

### Function

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed:<a href="http://www.uniprot.org/citations/1508712" target="\_blank">1508712</a>, PubMed:<a href="http://www.uniprot.org/citations/8183370" target="\_blank">8183370</a>). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed:<a href="http://www.uniprot.org/citations/11368354" target="\_blank">11368354</a>). Appears to function in a feedback loop system with caspases (PubMed:<a href="http://www.uniprot.org/citations/11368354" target="\_blank">11368354</a>). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed:<a href="http://www.uniprot.org/citations/11368354" target="\_blank">11368354</a>). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during non-starvation conditions and inhibits their autophagy function (PubMed:<a href="http://www.uniprot.org/citations/18570871" target="\_blank">18570871</a>, PubMed:<a href="http://www.uniprot.org/citations/20889974" target="\_blank">20889974</a>, PubMed:<a href="http://www.uniprot.org/citations/21358617" target="\_blank">21358617</a>). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:<a href="http://www.uniprot.org/citations/17418785" target="\_blank">17418785</a>).

### Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P10417}

### Tissue Location

Expressed in a variety of tissues.

## Anti-Bcl-2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Bcl-2 Picoband Antibody - Images

## Anti-Bcl-2 Picoband Antibody - Background

Immunoreactive BCL2 protein in the neoplastic cells of almost all follicular lymphomas whereas no BCL2 protein was detected in follicles affected by nonneoplastic processes or in normal lymphoid tissue. Every tumor with molecular-genetic evidence of t(14;18) translocation expressed detectable levels of BCL2 protein, regardless of whether the breakpoint was located in or at a distance from the BCL2 gene. Overexpression of BCL2 blocks the apoptotic death of a pro-B-lymphocyte cell line.