

Anti-PRKAR1A Picoband Antibody
Catalog # ABO10102**Specification****Anti-PRKAR1A Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	P10644
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for cAMP-dependent protein kinase type I-alpha regulatory subunit(PRKAR1A) detection. Tested with WB, IHC-P in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-PRKAR1A Picoband Antibody - Additional Information

Gene ID 5573

Other Names

cAMP-dependent protein kinase type I-alpha regulatory subunit, Tissue-specific extinguisher 1, TSE1, cAMP-dependent protein kinase type I-alpha regulatory subunit, N-terminally processed, PRKAR1A, PKR1, PRKAR1, TSE1

Calculated MW

42982 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Cell membrane .

Tissue Specificity

Four types of regulatory chains are found: I- alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible.

Protein Name

cAMP-dependent protein kinase type I-alpha regulatory subunit

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E.coli-derived human PRKAR1A recombinant protein (Position: E2-E81). Human PRKAR1A shares

89.9% and 92.4% amino acid (aa) sequence identity with mouse and rat PRKAR1A, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-PRKAR1A Picoband Antibody - Protein Information

Name PRKAR1A

Synonyms PKR1, PRKAR1, TSE1

Function

Regulatory subunit of the cAMP-dependent protein kinases involved in cAMP signaling in cells.

Cellular Location

Cell membrane.

Tissue Location


Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible


Anti-PRKAR1A Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PRKAR1A Picoband Antibody - Images







Western blot analysis of PRKAR1A expression in rat thymus extract (lane 1), HEPG-2 whole cell lysates (lane 2) and MCF-7 whole cell lysates (lane 3). PRKAR1A at 48KD was detected using rabbit anti- PRKAR1A Antigen Affinity purified polyclonal antibody (Catalog #ABO10102) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .



PRKAR1A was detected in paraffin-embedded sections of rat testis tissues using rabbit anti-PRKAR1A Antigen Affinity purified polyclonal antibody (Catalog # ABO10102) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



PRKAR1A was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- PRKAR1A Antigen Affinity purified polyclonal antibody (Catalog # ABO10102) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



PRKAR1A was detected in paraffin-embedded sections of human lung cancer tissues using rabbit anti- PRKAR1A Antigen Affinity purified polyclonal antibody (Catalog # ABO10102) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Anti-PRKAR1A Picoband Antibody - Background

cAMP-dependent protein kinase type I-alpha regulatory subunit is an enzyme that in humans is encoded by the PRKAR1A gene. This protein encoded by this gene was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids. Mutations in this gene cause Carney complex (CNC). This gene can fuse to the RET protooncogene by gene rearrangement and form the thyroid tumor-specific chimeric oncogene known as PTC2. A nonconventional nuclear localization sequence (NLS) has been found for this protein which suggests a role in DNA replication via the protein serving as a nuclear transport protein for the second subunit of the Replication Factor C (RFC40). Several alternatively spliced transcript variants encoding two different isoforms have been observed.