

**Anti-ETS1 Picoband Antibody**  
Catalog # ABO10127**Specification****Anti-ETS1 Picoband Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P14921</a>
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Protein C-ets-1(ETS1) detection. Tested with WB in Human;Mouse.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-ETS1 Picoband Antibody - Additional Information**

**Gene ID** 2113

**Other Names**

Protein C-ets-1, p54, ETS1, EWSR2

**Calculated MW**

50408 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Mouse<br>

**Subcellular Localization**

Cytoplasm . Nucleus . Delocalizes from nucleus to cytoplasm when coexpressed with isoform Ets-1 p27. .

**Tissue Specificity**

Highly expressed within lymphoid cells. Isoforms c-ETS-1A and Ets-1 p27 are both detected in all fetal tissues tested, but vary with tissue type in adult tissues. None is detected in brain or kidney. .

**Protein Name**

Protein C-ets-1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human ETS1 (67-98aa K DPRQWTETHVRDWVMWAVNEFSLKGVDFQKF), identical to the related mouse and rat sequences.

### Purification

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins.

### Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

## Anti-ETS1 Picoband Antibody - Protein Information

**Name** ETS1

**Synonyms** EWSR2

### Function

Transcription factor (PubMed:<a href="http://www.uniprot.org/citations/10698492" target="\_blank">10698492</a>, PubMed:<a href="http://www.uniprot.org/citations/11909962" target="\_blank">11909962</a>). Directly controls the expression of cytokine and chemokine genes in a wide variety of different cellular contexts (PubMed:<a href="http://www.uniprot.org/citations/20378371" target="\_blank">20378371</a>). May control the differentiation, survival and proliferation of lymphoid cells (PubMed:<a href="http://www.uniprot.org/citations/20378371" target="\_blank">20378371</a>). May also regulate angiogenesis through regulation of expression of genes controlling endothelial cell migration and invasion (PubMed:<a href="http://www.uniprot.org/citations/15247905" target="\_blank">15247905</a>, PubMed:<a href="http://www.uniprot.org/citations/15592518" target="\_blank">15592518</a>).

### Cellular Location

Nucleus. Cytoplasm Note=Delocalizes from nucleus to cytoplasm when coexpressed with isoform Ets-1 p27.

### Tissue Location


Highly expressed within lymphoid cells. Isoforms c- ETS-1A and Ets-1 p27 are both detected in all fetal tissues tested, but vary with tissue type in adult tissues. None is detected in brain or kidney.

## Anti-ETS1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-ETS1 Picoband Antibody - Images






Figure 1. Western blot analysis of ETS1 using anti- ETS1 antibody (ABO10127). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: NIH3T3 whole Cell lysates, Lane 2: A375 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti- ETS1 antigen affinity purified polyclonal antibody (Catalog # ABO10127) at 0.5  $\mu$ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ETS1 at approximately 54KD. The expected band size for ETS1 is at 54KD.

### Anti-ETS1 Picoband Antibody - Background

Protein C-ets-1 is a protein that in humans is encoded by the ETS1 gene. It is mapped to 11q24.3. This gene encodes a member of the ETS family of transcription factors, which are defined by the presence of a conserved ETS DNA-binding domain that recognizes the core consensus DNA sequence GGAA/T in target genes. These proteins function either as transcriptional activators or repressors of numerous genes, and are involved in stem cell development, cell senescence and death, and tumorigenesis.