

Anti-IRF9 Picoband Antibody
Catalog # ABO10307**Specification****Anti-IRF9 Picoband Antibody - Product Information**

Application	WB
Primary Accession	Q00978
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Interferon regulatory factor 9 (IRF9) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IRF9 Picoband Antibody - Additional Information

Gene ID 10379

Other Names

Interferon regulatory factor 9, IRF-9, IFN-alpha-responsive transcription factor subunit, ISGF3 p48 subunit, Interferon-stimulated gene factor 3 gamma, ISGF-3 gamma, Transcriptional regulator ISGF3 subunit gamma, IRF9, ISGF3G

Calculated MW

43696 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Cytoplasm . Nucleus . Translocated into the nucleus upon activation by IFN-alpha/beta.

Protein Name

Interferon regulatory factor 9

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E. coli-derived human IRF9 recombinant protein (Position: M1-K110). Human IRF9 shares 87.3% amino acid (aa) sequence identity with mouse IRF9.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-IRF9 Picoband Antibody - Protein Information

Name IRF9

Synonyms ISGF3G

Function

Transcription factor that plays an essential role in anti-viral immunity. It mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. IRF9/ISGF3G associates with the phosphorylated STAT1:STAT2 dimer to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state.


Cellular Location


Cytoplasm. Nucleus Note=Translocated into the nucleus upon activation by IFN-alpha/beta

Anti-IRF9 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IRF9 Picoband Antibody - Images



Western blot analysis of IRF9 expression in U87 whole cell lysates (lane 1). IRF9 at 44KD was detected using rabbit anti- IRF9 Antigen Affinity purified polyclonal antibody (Catalog # ABO10307) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-IRF9 Picoband Antibody - Background

Interferon regulatory factor 9 is a protein that in humans is encoded by the IRF9 gene, previously known as ISGF3G. It is mapped to 14q1. IRF9 is a 48 kDa member of the IRF family of proteins. It is widely expressed, and serves as a component of the ISGF3 complex. In addition, IRF9 is an interferon-dependent, positive-acting transcription factor that is cytoplasmically activated, possibly through direct interaction with the interferon receptor. It associates with activated STAT1 and STAT2 to form an ISGF3 complex that is translocated into the nucleus.