

Anti-BNIP3 Antibody
Catalog # ABO10543**Specification****Anti-BNIP3 Antibody - Product Information**

Application	WB
Primary Accession	Q12983
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for BCL2/adenovirus E1B 19 kDa protein-interacting protein 3(BNIP3) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-BNIP3 Antibody - Additional Information

Gene ID 664

Other Names

BCL2/adenovirus E1B 19 kDa protein-interacting protein 3, BNIP3, NIP3

Calculated MW

21541 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Mitochondrion. Mitochondrion outer membrane; Single-pass membrane protein. Coexpression with the E1B 19- kDa protein results in a shift in NIP3 localization pattern to the nuclear envelope. Colocalizes with ACAA2 in the mitochondria. Colocalizes with SPATA18 at the mitochondrion outer membrane.

Protein Name

BCL2/adenovirus E1B 19 kDa protein-interacting protein 3

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human BNIP3(38-53aa IYNGDMEKILLDAQHE), identical to the related mouse and rat sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the NIP3 family.

Anti-BNIP3 Antibody - Protein Information

Name BNIP3 ([HGNC:1084](#))

Synonyms NIP3

Function

Apoptosis-inducing protein that can overcome BCL2 suppression. May play a role in repartitioning calcium between the two major intracellular calcium stores in association with BCL2. Involved in mitochondrial quality control via its interaction with SPATA18/MIEAP: in response to mitochondrial damage, participates in mitochondrial protein catabolic process (also named MALM) leading to the degradation of damaged proteins inside mitochondria. The physical interaction of SPATA18/MIEAP, BNIP3 and BNIP3L/NIX at the mitochondrial outer membrane regulates the opening of a pore in the mitochondrial double membrane in order to mediate the translocation of lysosomal proteins from the cytoplasm to the mitochondrial matrix. Plays an important role in the calprotectin (S100A8/A9)-induced cell death pathway.

Cellular Location

Mitochondrion. Mitochondrion outer membrane; Single-pass membrane protein.


Note=Coexpression with the E1B 19-kDa protein results in a shift in NIP3 localization pattern to the nuclear envelope. Colocalizes with ACAA2 in the mitochondria. Colocalizes with SPATA18 at the mitochondrion outer membrane


Anti-BNIP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-BNIP3 Antibody - Images





Anti-BNIP3 antibody, ABO10543, Western blotting All lanes: Anti BNIP3 (ABO10543) at 0.5ug/ml Lane 1: HELA Whole Cell Lysate at 40ug Lane 2: MM231 Whole Cell Lysate at 40ug Predicted bind size: 22KD Observed bind size: 30KD

Anti-BNIP3 Antibody - Background

The Bcl-2 nineteen kilodalton interacting protein 3 (BNIP3 or NIP3), is a hypoxia-inducible proapoptotic member of the Bcl-2 family that induces cell death by associating with the mitochondria. BNIP3, expressed in skeletal muscle and in the brain at low levels, is primarily localized to the nucleus of glial cells of the normal human brain, as well as in the malignant glioma cell line U251. BNIP3 expression in the cytoplasm increases and localizes with the mitochondria, contributing to induction of cell death. Cellular protein BNIP3 interacts with E1B-19K, BCL-2, BCL-xL, and EBV-BHRF1. BNIP3 contains Bcl-2 homology 3 (BH3) domain and COOH-terminal transmembrane (TM) domain. The BH3 domain of BNIP3 mediates Bcl-2/Bcl-X(L) heterodimerization and confers pro-apoptotic activity; whereas the TM domain is critical for homodimerization, pro-apoptotic function, and mitochondrial targeting.