

**Anti-STAT1 Antibody**  
Catalog # ABO10558

**Specification**

---

**Anti-STAT1 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P42224</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Signal transducer and activator of transcription 1-alpha/beta(STAT1) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-STAT1 Antibody - Additional Information**

**Gene ID** 6772

**Other Names**

Signal transducer and activator of transcription 1-alpha/beta, Transcription factor ISGF-3 components p91/p84, STAT1

**Calculated MW**

87335 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat<br>

**Subcellular Localization**

Cytoplasm . Nucleus . Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to IFN-gamma and signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4.

**Protein Name**

Signal transducer and activator of transcription 1-alpha/beta

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human STAT1(364-378aa FDKDVNERNTVKGFR), different from the related mouse sequence by one amino acid.

**Purification**

Immunogen affinity purified.

## Cross Reactivity

No cross reactivity with other proteins

## Storage

**At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

## Sequence Similarities

Belongs to the transcription factor STAT family.

## Anti-STAT1 Antibody - Protein Information

### Name STAT1

### Function

Signal transducer and transcription activator that mediates cellular responses to interferons (IFNs), cytokine KITLG/SCF and other cytokines and other growth factors (PubMed:<a href="http://www.uniprot.org/citations/12764129" target="\_blank">12764129</a>, PubMed:<a href="http://www.uniprot.org/citations/12855578" target="\_blank">12855578</a>, PubMed:<a href="http://www.uniprot.org/citations/15322115" target="\_blank">15322115</a>, PubMed:<a href="http://www.uniprot.org/citations/23940278" target="\_blank">23940278</a>, PubMed:<a href="http://www.uniprot.org/citations/34508746" target="\_blank">34508746</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>, PubMed:<a href="http://www.uniprot.org/citations/9724754" target="\_blank">9724754</a>). Following type I IFN (IFN-alpha and IFN-beta) binding to cell surface receptors, signaling via protein kinases leads to activation of Jak kinases (TYK2 and JAK1) and to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize and associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus (PubMed:<a href="http://www.uniprot.org/citations/28753426" target="\_blank">28753426</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>). ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IFN-stimulated genes (ISG), which drive the cell in an antiviral state (PubMed:<a href="http://www.uniprot.org/citations/28753426" target="\_blank">28753426</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>). In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated (PubMed:<a href="http://www.uniprot.org/citations/26479788" target="\_blank">26479788</a>). It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state (PubMed:<a href="http://www.uniprot.org/citations/8156998" target="\_blank">8156998</a>). Becomes activated in response to KITLG/SCF and KIT signaling (PubMed:<a href="http://www.uniprot.org/citations/15526160" target="\_blank">15526160</a>). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed:<a href="http://www.uniprot.org/citations/19088846" target="\_blank">19088846</a>). Following bacterial lipopolysaccharide (LPS)-induced TLR4 endocytosis, phosphorylated at Thr-749 by IKBKB which promotes binding of STAT1 to the 5'-TTTGAGGC-3' sequence in the ARID5A promoter, resulting in transcriptional activation of ARID5A and subsequent ARID5A-mediated stabilization of IL6 (PubMed:<a href="http://www.uniprot.org/citations/32209697" target="\_blank">32209697</a>). Phosphorylation at Thr-749 also promotes binding of STAT1 to the 5'-TTTGAGTC-3' sequence in the IL12B promoter and activation of IL12B transcription (PubMed:<a href="http://www.uniprot.org/citations/32209697" target="\_blank">32209697</a>). Involved in food tolerance in small intestine: associates with the Gasdermin-D, p13 cleavage product (13 kDa GSDMD) and promotes transcription of CIITA, inducing type 1 regulatory T (Tr1) cells in upper small intestine (By similarity).

### Cellular Location


Cytoplasm. Nucleus Note=Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to IFN-gamma and signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4 (PubMed:15322115). Monomethylation at Lys- 525 is required for phosphorylation at Tyr-701 and translocation into the nucleus (PubMed:28753426). Translocates into the nucleus in response to interferon-beta stimulation (PubMed:26479788)

### Anti-STAT1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-STAT1 Antibody - Images



Anti-STAT1 antibody, ABO10558, Western blotting Lane 1: MCF-7 Cell Lysate Lane 2: HELA Cell Lysate



Anti-STAT1 antibody, ABO10558, Western blotting  
Lane 1: Mouse Heart Tissue Lysate  
Lane 2: Mouse Liver Tissue Lysate  
Lane 3: Mouse Brain Tissue Lysate  
Lane 4: Mouse Kidney Tissue Lysate  
Lane 5: Mouse Spleen Tissue Lysate  
Lane 6: Mouse Thymus Tissue Lysate  
Lane 7: Mouse Lung Tissue Lysate  
Lane 8: Mouse Intestine Tissue Lysate  
Lane 9: Mouse Ovary Tissue Lysate

### **Anti-STAT1 Antibody - Background**

Signal transducer and activator of transcription 1 (STAT1) is a transcription factor which in humans is encoded by the STAT1 gene. The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described.