

Anti-GFAP Antibody
Catalog # ABO10614**Specification**

Anti-GFAP Antibody - Product Information

Application	WB, IHC
Primary Accession	P14136
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Glial fibrillary acidic protein(GFAP) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-GFAP Antibody - Additional Information

Gene ID 2670

Other Names

Glial fibrillary acidic protein, GFAP, GFAP

Calculated MW

49880 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Rat, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cytoplasm . Associated with intermediate filaments.

Tissue Specificity

Expressed in cells lacking fibronectin. .

Protein Name

Glial fibrillary acidic protein(GFAP)

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human GFAP(417-432aa DGEVIKESKQEHKDVM), identical to the related rat sequence, and different from the related mouse sequence by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the intermediate filament family.

Anti-GFAP Antibody - Protein Information

Name GFAP

Function

GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.

Cellular Location

Cytoplasm. Note=Associated with intermediate filaments


Tissue Location


Expressed in cells lacking fibronectin.

Anti-GFAP Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-GFAP Antibody - Images



Anti-GFAP antibody, ABO10614, Western blotting All lanes: Anti GFAP(ABO10614) at 0.5ug/ml
 Lane 1: Rat Brain Tissue Lysate at 50ug
 Lane 2: Mouse Brain Tissue Lysate at 50ug
 Lane 3: U87 Whole Cell Lysate at 40ug
 Predicted bind size: 49KD
 Observed bind size: 49KD



Anti-GFAP antibody, ABO10614, IHC(P)
 IHC(P): Rat Brain Tissue



Anti-GFAP antibody, ABO10614, Western blotting
 Lane 1: Rat Brain Tissue Lysate
 Lane 2: Rat Brain Tissue Lysate
 Lane 3: Mouse Brain Tissue Lysate
 Lane 4: Mouse Brain Tissue Lysate

Anti-GFAP Antibody - Background

Glial fibrillary acidic protein(GFAP) is an intermediate-filament(IF) protein that is highly specific for cells of astroglial lineage, although its tissue-specific role is speculative. GFAP has been located in rat kidney glomeruli and peritubular fibroblasts, leydig cells of testis, skin keratinocytes, osteocytes of bones, chondrocytes of epiglottis, bronchus, and stellate-shaped cells of the pancreas and liver. Its expression is essential for normal white matter architecture and blood-brain barrier integrity, and its absence leads to late-onset CNS dysmyelination. GFAP has also been shown to play a role in mitosis by adjusting the filament network present in the cell. During mitosis, there is an increase in the amount of phosphorylated GFAP, and a movement of this modified protein to the cleavage furrow.