

Anti-sTNFsR II Antibody
Catalog # ABO10617**Specification****Anti-sTNFsR II Antibody - Product Information**

Application	WB, IHC
Primary Accession	P20333
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Tumor necrosis factor receptor superfamily member 1B(TNFRSF1B) detection. Tested with WB, IHC-P in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-sTNFsR II Antibody - Additional Information

Gene ID 7133

Other Names

Tumor necrosis factor receptor superfamily member 1B, Tumor necrosis factor receptor 2, TNF-R2, Tumor necrosis factor receptor type II, TNF-RII, TNFR-II, p75, p80 TNF-alpha receptor, CD120b, Etanercept, Tumor necrosis factor receptor superfamily member 1b, membrane form, Tumor necrosis factor-binding protein 2, TBP-2, TBPII, TNFRSF1B, TNFBR, TNFR2

Calculated MW

48291 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Isoform 1: Cell membrane; Single-pass type I membrane protein.

Protein Name

Tumor necrosis factor receptor superfamily member 1B

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg Na₂S₂O₃.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human TNF Receptor II(103-121aa DQVETQACTREQNRICTCR).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Contains 4 TNFR-Cys repeats.

Anti-sTNFsR II Antibody - Protein Information

Name TNFRSF1B

Synonyms TNFBR, TNFR2

Function

Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.


Cellular Location


[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Tumor necrosis factor-binding protein 2]: Secreted

Anti-sTNFsR II Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-sTNFsR II Antibody - Images



Anti-TNF Receptor II antibody, ABO10617, Western blotting Lane 1: MM453 Cell Lysate Lane 2: JURKAT Cell Lysate



Anti-TNF Receptor II antibody, ABO10617, IHC(P)IHC(P): Human Mammary Cancer Tissue

Anti-sTNFsR II Antibody - Background

Tumor necrosis factor receptor 2 (TNFR2) is one of the receptors of TNF. TNF has proinflammatory and immunosuppressive properties that may segregate at the level of the 2 TNF receptors (TNFRs), TNFR1 and TNFR2. The genes for TNFR1, a 55-kDa protein, and TNFR2, a 70-kDa protein, have been mapped to human chromosomes 12(12pter-cen) and 1(1pter-p32), respectively. TNFR2 was induced on glomerular endothelial cells of nephritic kidneys, and TNFR2 expression on intrinsic cells, but not leukocytes, was essential for glomerulonephritis and glomerular complement deposition. TNFR1 promotes systemic immune responses and renal T cell death, while intrinsic cell TNFR2 plays a critical role in complement-dependent tissue injury. Therefore, therapeutic blockade specifically of TNFR2 may be a promising strategy in the treatment of immune-mediated glomerulonephritis.