

**Anti-LXR Alpha Antibody**  
Catalog # ABO10622**Specification**

---

**Anti-LXR Alpha Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q13133</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Oxysterols receptor LXR-alpha(NR1H3) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-LXR Alpha Antibody - Additional Information**

**Gene ID** 10062

**Other Names**

Oxysterols receptor LXR-alpha, Liver X receptor alpha, Nuclear receptor subfamily 1 group H member 3, NR1H3, LXRA

**Calculated MW**

50396 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse<br>

**Subcellular Localization**

Nucleus .

**Tissue Specificity**

Visceral organs specific expression. Strong expression was found in liver, kidney and intestine followed by spleen and to a lesser extent the adrenals.

**Protein Name**

Oxysterols receptor LXR-alpha

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human LXR alpha(371-385aa NVQDQLQVERLQHTY), identical to the related rat and mouse sequences.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the nuclear hormone receptor family. NR1 subfamily.

**Anti-LXR Alpha Antibody - Protein Information**

**Name** NR1H3

**Synonyms** LXRA

**Function**

Nuclear receptor that exhibits a ligand-dependent transcriptional activation activity (PubMed:<a href="http://www.uniprot.org/citations/19481530" target="\_blank">19481530</a>, PubMed:<a href="http://www.uniprot.org/citations/25661920" target="\_blank">25661920</a>). Interaction with retinoic acid receptor (RXR) shifts RXR from its role as a silent DNA-binding partner to an active ligand-binding subunit in mediating retinoid responses through target genes defined by LXRES (By similarity). LXRES are DR4-type response elements characterized by direct repeats of two similar hexanuclotide half-sites spaced by four nucleotides (By similarity). Plays an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8 (PubMed:<a href="http://www.uniprot.org/citations/19481530" target="\_blank">19481530</a>). Interplays functionally with RORA for the regulation of genes involved in liver metabolism (By similarity). Induces LPCAT3- dependent phospholipid remodeling in endoplasmic reticulum (ER) membranes of hepatocytes, driving SREBF1 processing and lipogenesis (By similarity). Via LPCAT3, triggers the incorporation of arachidonate into phosphatidylcholines of ER membranes, increasing membrane dynamics and enabling triacylglycerols transfer to nascent very low-density lipoprotein (VLDL) particles. Via LPCAT3 also counteracts lipid-induced ER stress response and inflammation, likely by modulating SRC kinase membrane compartmentalization and limiting the synthesis of lipid inflammatory mediators (By similarity).

**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:25661920}.  
Cytoplasm {ECO:0000250|UniProtKB:Q9Z0Y9}

**Tissue Location**

Visceral organs specific expression. Strong expression was found in liver, kidney and intestine followed by spleen and to a lesser extent the adrenals

**Anti-LXR Alpha Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-LXR Alpha Antibody - Images



Anti- NR1H3 antibody, ABO10622, Western blotting All lanes: Anti NR1H3 (ABO10622) at 0.5ug/ml  
Lane 1: Rat Liver Tissue Lysate at 50ug  
Lane 2: Rat Lung Tissue Lysate at 50ug  
Lane 3: Rat Spleen Tissue Lysate at 50ug  
Lane 4: Rat Kidney Tissue Lysate at 50ug  
Lane 5: Human Placenta Tissue Lysate at 50ug  
Predicted bind size: 51KD  
Observed bind size: 51KD

### Anti-LXR Alpha Antibody - Background

LXRA is a tissue-specific cofactor that permits RXRA to function as a potent 9cRA receptor with a distinct target gene specificity. It specifically interacts with RXRA in vivo to form a functional heterodimer in which RXRA is the ligand-binding subunit. Additionally, LXR activity is critical for physiologic lipid metabolism and transport. LXRs are endogenous inhibitors of atherogenesis and are targets for therapeutic intervention in cardiovascular disease. Furthermore, LXRs and their ligands are negative regulators of macrophage inflammatory gene expression. LXR is also found that as a transcriptional switch that integrates hepatic glucose metabolism and fatty acid synthesis. The LXR-IDOL-LDLR axis defines a complementary pathway to sterol response element-binding proteins for sterol regulation of cholesterol uptake.