

Anti-VEGFD Antibody
Catalog # ABO10658

Specification

Anti-VEGFD Antibody - Product Information

Application	WB
Primary Accession	O43915
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Vascular endothelial growth factor D(FIGF) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-VEGFD Antibody - Additional Information

Gene ID 2277

Other Names

Vascular endothelial growth factor D {ECO:0000312|HGNC:HGNC:3708}, VEGF-D, c-Fos-induced growth factor, FIGF, VEGFD (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3708)>HGNC:3708), FIGF

Calculated MW

40444 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Secreted.

Tissue Specificity

Highly expressed in lung, heart, small intestine and fetal lung, and at lower levels in skeletal muscle, colon, and pancreas.

Protein Name

Vascular endothelial growth factor D

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human VEGFD(101-115aa VIDEEWQRTQCSPRE), identical to the related rat and mouse sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the PDGF/VEGF growth factor family.

Anti-VEGFD Antibody - Protein Information

Name VEGFD ([HGNC:3708](#))

Synonyms FIGF

Function

Growth factor active in angiogenesis, lymphangiogenesis and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in the formation of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates VEGFR-2 (KDR/FLK1) and VEGFR-3 (FLT4) receptors.

Cellular Location

Secreted.

Tissue Location


Highly expressed in lung, heart, small intestine and fetal lung, and at lower levels in skeletal muscle, colon, and pancreas


Anti-VEGFD Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-VEGFD Antibody - Images





Anti-VEGFD antibody, ABO10658, Western blotting Lane 1: SW620 Cell Lysate Lane 2: COLO320 Cell Lysate Lane 3: 6T-CEM Cell Lysate Lane 4: HT1080 Cell Lysate

Anti-VEGFD Antibody - Background

C-fos induced growth factor (FIGF) (or vascular endothelial growth factor D, VEGF-D) is a vascular endothelial growth factor that in humans is encoded by the FIGF gene. The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family and is active in angiogenesis, lymphangiogenesis, and endothelial cell growth. Analyzing by Northern blotting, Yamada et al. (1997) symbolized VEGFD, was expressed as a 2.2-kb transcript with highest expression in lung, heart, small intestine, and fetal lung, and lower levels in skeletal muscle, colon, and pancreas. And Achen et al. (1998) concluded that VEGFD was most closely related to VEGFC by virtue of the presence of N- and C-terminal extensions that were not found in other VEGF family members. Stacker et al. (2001) showed that VEGFD can induce tumor angiogenesis through VEGFR2 and tumor lymphangiogenesis through VEGFR3, whereas VEGF, which does not activate VEGFR3, induces only tumor angiogenesis.