

**Anti-IL-10 Antibody**  
Catalog # ABO10678**Specification**

---

**Anti-IL-10 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P22301</a>
Host	<b>Rabbit</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Format	<b>Lyophilized</b>

**Description**

Rabbit IgG polyclonal antibody for Interleukin-10(IL10) detection. Tested with WB in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-IL-10 Antibody - Additional Information**

**Gene ID** 3586

**Other Names**

Interleukin-10, IL-10, Cytokine synthesis inhibitory factor, CSIF, IL10

**Calculated MW**

20517 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted.

**Tissue Specificity**

Produced by a variety of cell lines, including T-cells, macrophages, mast cells and other cell types.

**Protein Name**

Interleukin-10(IL-10)

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human IL10(46-60aa DAFSRVKTFQMKDQ).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the IL-10 family.

**Anti-IL-10 Antibody - Protein Information**

**Name** IL10

**Function**

Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3 (PubMed: [16982608](http://www.uniprot.org/citations/16982608)). In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators (PubMed: [18025162](http://www.uniprot.org/citations/18025162)). Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro-inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha (PubMed: [11564774](http://www.uniprot.org/citations/11564774), PubMed: [1940799](http://www.uniprot.org/citations/1940799), PubMed: [7512027](http://www.uniprot.org/citations/7512027)). Interferes also with antigen presentation by reducing the expression of MHC-class II and co-stimulatory molecules, thereby inhibiting their ability to induce T cell activation (PubMed: [8144879](http://www.uniprot.org/citations/8144879)). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).

**Cellular Location**

Secreted.

**Tissue Location**


Produced by a variety of cell lines, including T- cells, macrophages, mast cells and other cell types

**Anti-IL-10 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-IL-10 Antibody - Images



Anti-IL-10 antibody, ABO10678, Western blotting Lane 1: Recombinant Human IL-10 Protein 10ng Lane 2: Recombinant Human IL-10 Protein 5ng Lane 3: Recombinant Human IL-10 Protein 2.5ng

## Anti-IL-10 Antibody - Background

Interleukin-10(IL-10 or IL10), also known as human cytokine synthesis inhibitory factor(CSIF), is an anti-inflammatory cytokine. In humans IL-10 is encoded by the IL10 gene. It is capable of inhibiting synthesis of pro-inflammatory cytokines like IFN-gamma, IL-2, IL-3, TNFalpha and GM-CSF made by cells such as macrophages and regulatory T-cells. IL-10 also displays potent abilities to suppress the antigen presentation capacity of antigen presenting cells. Kim et al.(1992) showed that the mouse IL 10 gene contains 5 exons and spans about 5.2 kb of genomic DNA. Eskdale et al.(1997) mapped the IL10 gene to the junction between 1q31 and 1q32.