

Anti-Midkine Antibody
Catalog # ABO10848**Specification**

Anti-Midkine Antibody - Product Information

Application	WB
Primary Accession	P21741
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Midkine(MDK) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Midkine Antibody - Additional Information

Gene ID 4192

Other Names

Midkine, MK, Amphiregulin-associated protein, ARAP, Midgestation and kidney protein, Neurite outgrowth-promoting factor 2, Neurite outgrowth-promoting protein, MDK, MK1, NEGF2

Calculated MW

15585 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Secreted.

Tissue Specificity

Expressed in various tumor cell lines. In insulinoma tissue predominantly expressed in precancerous lesions. .

Protein Name

Midkine(MK)

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Midkine(108-121aa KKARYNAQCQETIR), identical to the related rat and mouse sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-Midkine Antibody - Protein Information

Name MDK ([HGNC:6972](#))

Synonyms MK1, NRG2

Function

Secreted protein that functions as a cytokine and growth factor and mediates its signal through cell-surface proteoglycan and non-proteoglycan receptors (PubMed:[10212223](http://www.uniprot.org/citations/10212223), PubMed:[10772929](http://www.uniprot.org/citations/10772929), PubMed:[12084985](http://www.uniprot.org/citations/12084985), PubMed:[12122009](http://www.uniprot.org/citations/12122009), PubMed:[12573468](http://www.uniprot.org/citations/12573468), PubMed:[15466886](http://www.uniprot.org/citations/15466886), PubMed:[18469519](http://www.uniprot.org/citations/18469519), PubMed:[24458438](http://www.uniprot.org/citations/24458438)). Binds cell-surface proteoglycan receptors via their chondroitin sulfate (CS) groups (PubMed:[10212223](http://www.uniprot.org/citations/10212223), PubMed:[12084985](http://www.uniprot.org/citations/12084985)). Thereby regulates many processes like inflammatory response, cell proliferation, cell adhesion, cell growth, cell survival, tissue regeneration, cell differentiation and cell migration (PubMed:[10212223](http://www.uniprot.org/citations/10212223), PubMed:[10683378](http://www.uniprot.org/citations/10683378), PubMed:[10772929](http://www.uniprot.org/citations/10772929), PubMed:[12084985](http://www.uniprot.org/citations/12084985), PubMed:[12122009](http://www.uniprot.org/citations/12122009), PubMed:[12573468](http://www.uniprot.org/citations/12573468), PubMed:[15466886](http://www.uniprot.org/citations/15466886), PubMed:[22323540](http://www.uniprot.org/citations/22323540), PubMed:[24458438](http://www.uniprot.org/citations/24458438)). Participates in inflammatory processes by exerting two different activities. Firstly, mediates neutrophils and macrophages recruitment to the sites of inflammation both by direct action by cooperating namely with ITGB2 via LRP1 and by inducing chemokine expression (PubMed:[10683378](http://www.uniprot.org/citations/10683378), PubMed:[24458438](http://www.uniprot.org/citations/24458438)). This inflammation can be accompanied by epithelial cell survival and smooth muscle cell migration after renal and vessel damage, respectively (PubMed:[10683378](http://www.uniprot.org/citations/10683378)). Secondly, suppresses the development of tolerogenic dendric cells thereby inhibiting the differentiation of regulatory T cells and also promote T cell expansion through NFAT signaling and Th1 cell differentiation (PubMed:[22323540](http://www.uniprot.org/citations/22323540)). Promotes tissue regeneration after injury or trauma. After heart damage negatively regulates the recruitment of inflammatory cells and mediates cell survival through activation of anti-apoptotic signaling pathways via MAPKs and AKT pathways through the

activation of angiogenesis (By similarity). Also facilitates liver regeneration as well as bone repair by recruiting macrophage at trauma site and by promoting cartilage development by facilitating chondrocyte differentiation (By similarity). Plays a role in brain by promoting neural precursor cells survival and growth through interaction with heparan sulfate proteoglycans (By similarity). Binds PTPRZ1 and promotes neuronal migration and embryonic neurons survival (PubMed:10212223). Binds SDC3 or GPC2 and mediates neurite outgrowth and cell adhesion (PubMed:12084985, PubMed:1768439). Binds chondroitin sulfate E and heparin leading to inhibition of neuronal cell adhesion induced by binding with GPC2 (PubMed:12084985). Binds CSPG5 and promotes elongation of oligodendroglial precursor-like cells (By similarity). Also binds ITGA6:ITGB1 complex; this interaction mediates MDK-induced neurite outgrowth (PubMed:15466886, PubMed:1768439). Binds LRP1; promotes neuronal survival (PubMed:10772929). Binds ITGA4:ITGB1 complex; this interaction mediates MDK-induced osteoblast cells migration through PXN phosphorylation (PubMed:15466886). Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated protein kinase (MAPK) and PI3-kinase, and the induction of cell proliferation (PubMed:12122009). Promotes epithelial to mesenchymal transition through interaction with NOTCH2 (PubMed:18469519). During arteriogenesis, plays a role in vascular endothelial cell proliferation by inducing VEGFA expression and release which in turn induces nitric oxide synthase expression. Moreover activates vasodilation through nitric oxide synthase activation (By similarity). Negatively regulates bone formation in response to mechanical load by inhibiting Wnt/beta- catenin signaling in osteoblasts (By similarity). In addition plays a role in hippocampal development, working memory, auditory response, early fetal adrenal gland development and the female reproductive system (By similarity).

Cellular Location

Secreted.

Tissue Location

Expressed in various tumor cell lines. In insulinoma tissue predominantly expressed in precancerous lesions


Anti-Midkine Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Midkine Antibody - Images





Anti-Midkine Antibody - Background

Midkine(MK or MDK) also known as neurite growth-promoting factor 2(NEGF2) is a protein that in humans is encoded by the MDK gene. Midkine is a basic heparin-binding growth factor of low molecular weight, and forms a family with pleiotrophin(NEGF1, 46% homologous with MK). It is a nonglycosylated protein, composed of two domains held by disulfide bridges. It is a developmentally important retinoic acid-responsive gene product strongly induced during mid-gestation, hence the name midkine. Restricted mainly to certain tissues in the normal adult, it is strongly induced during oncogenesis, inflammation and tissue repair. MK is pleiotropic, capable of exerting activities such as cell proliferation, cell migration, angiogenesis and fibrinolysis. A molecular complex containing receptor-type tyrosine phosphatase zeta(PTP zeta), low density lipoprotein receptor-related protein(LRP1), anaplastic leukemia kinase(ALK) and syndecans is considered to be its receptor