

**Anti-Cdc25B Antibody**  
Catalog # ABO10863**Specification**

---

**Anti-Cdc25B Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P30305</a>
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for M-phase inducer phosphatase 2(CDC25B) detection. Tested with WB, IHC-P in Human;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Cdc25B Antibody - Additional Information**

**Gene ID** 994

**Other Names**

M-phase inducer phosphatase 2, 3.1.3.48, Dual specificity phosphatase Cdc25B, CDC25B, CDC25HU2

**Calculated MW**

64987 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, By Heat  
Western blot, 0.1-0.5 µg/ml, Rat, Human

**Subcellular Localization**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole.

**Protein Name**

M-phase inducer phosphatase 2

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human Cdc25B(541-559aa DYRPMNHEAFKDELKTFRL), different from the related mouse sequence by three amino acids.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### Sequence Similarities

Belongs to the MPI phosphatase family.

## Anti-Cdc25B Antibody - Protein Information

**Name** CDC25B

**Synonyms** CDC25HU2

### Function

Tyrosine protein phosphatase which functions as a dosage- dependent inducer of mitotic progression (PubMed: [1836978](http://www.uniprot.org/citations/1836978), PubMed: [20360007](http://www.uniprot.org/citations/20360007) target="\_blank">20360007</a>). Directly dephosphorylates CDK1 and stimulates its kinase activity (PubMed: [20360007](http://www.uniprot.org/citations/20360007) target="\_blank">20360007</a>). Required for G2/M phases of the cell cycle progression and abscission during cytokinesis in a ECT2-dependent manner (PubMed: [17332740](http://www.uniprot.org/citations/17332740) target="\_blank">17332740</a>). The three isoforms seem to have a different level of activity (PubMed: [1836978](http://www.uniprot.org/citations/1836978) target="\_blank">1836978</a>).

### Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole


## Anti-Cdc25B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Cdc25B Antibody - Images





Anti-Cdc25B antibody, ABO10863, Western blotting Lane 1: Rat Brain Tissue Lysate Lane 2: Rat Kidney Tissue Lysate Lane 3: Rat Lung Tissue Lysate



Anti-Cdc25B antibody, ABO10863, IHC(P) IHC(P): Human Lung Cancer Tissue

### **Anti-Cdc25B Antibody - Background**

Central to the onset of mitosis in all eukaryotic cells is the CDC2 protein kinase, the activity of which is negatively regulated by phosphorylation and positively activated by dephosphorylation. The latter function is carried out by a specific phosphatase, CDC25. At least 3 human CDC25 genes code for the A, B, and C forms of CDC25. CDC25B is mapped to 20p13. P38 kinase has a critical role in the initiation of a G2 delay after ultraviolet radiation. Inhibition of p38 blocks the rapid initiation of this checkpoint in both human and murine cells after ultraviolet radiation. In vitro, p38 binds and phosphorylates CDC25B at serines 309 and 361, and CDC25C at serine-216; phosphorylation of these residues is required for binding to 14-3-3 proteins. In vivo, inhibition of p38 prevents both phosphorylation of CDC25B at serine-309 and 14-3-3 binding after ultraviolet radiation, and mutation of this site is sufficient to inhibit the checkpoint initiation. Regulation of CDC25B phosphorylation by p38 is a critical event for initiating the G2/M checkpoint after ultraviolet radiation.