

**Anti-ALOX5 Antibody**  
Catalog # ABO10959

**Specification**

---

**Anti-ALOX5 Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P09917</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Arachidonate 5-lipoxygenase(ALOX5) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-ALOX5 Antibody - Additional Information**

**Gene ID** 240

**Other Names**

Arachidonate 5-lipoxygenase, 5-LO, 5-lipoxygenase, 1.13.11.34, ALOX5, LOG5

**Calculated MW**

77983 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br><br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Cytoplasm. Nucleus matrix. Nucleus membrane; Peripheral membrane protein. Shuttles between cytoplasm and nucleus. Found exclusively in the nucleus, when phosphorylated on Ser-272. Calcium binding promotes translocation from the cytosol and the nuclear matrix to the nuclear envelope and membrane association.

**Protein Name**

Arachidonate 5-lipoxygenase(5-LO/5-lipoxygenase)

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human ALOX5 (650-667aa AERNKKKQLPYYYYLSPDR), different from the related rat and mouse sequences by two amino acids.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

### Sequence Similarities

Belongs to the lipoxygenase family.

## Anti-ALOX5 Antibody - Protein Information

Name ALOX5 ([HGNC:435](#))

Synonyms LOG5

### Function

Catalyzes the oxygenation of arachidonate ((5Z,8Z,11Z,14Z)- eicosatetraenoate) to 5-hydroperoxyeicosatetraenoate (5-HPETE) followed by the dehydration to 5,6-epoxyeicosatetraenoate (Leukotriene A4/LTA4), the first two steps in the biosynthesis of leukotrienes, which are potent mediators of inflammation (PubMed:<a href="http://www.uniprot.org/citations/19022417" target="\_blank">19022417</a>, PubMed:<a href="http://www.uniprot.org/citations/21233389" target="\_blank">21233389</a>, PubMed:<a href="http://www.uniprot.org/citations/22516296" target="\_blank">22516296</a>, PubMed:<a href="http://www.uniprot.org/citations/23246375" target="\_blank">23246375</a>, PubMed:<a href="http://www.uniprot.org/citations/24282679" target="\_blank">24282679</a>, PubMed:<a href="http://www.uniprot.org/citations/24893149" target="\_blank">24893149</a>, PubMed:<a href="http://www.uniprot.org/citations/31664810" target="\_blank">31664810</a>, PubMed:<a href="http://www.uniprot.org/citations/8615788" target="\_blank">8615788</a>, PubMed:<a href="http://www.uniprot.org/citations/8631361" target="\_blank">8631361</a>). Also catalyzes the oxygenation of arachidonate into 8- hydroperoxyeicosatetraenoate (8-HPETE) and 12- hydroperoxyicosatetraenoate (12-HPETE) (PubMed:<a href="http://www.uniprot.org/citations/23246375" target="\_blank">23246375</a>). Displays lipoxin synthase activity being able to convert (15S)-HETE into a conjugate tetraene (PubMed:<a href="http://www.uniprot.org/citations/31664810" target="\_blank">31664810</a>). Although arachidonate is the preferred substrate, this enzyme can also metabolize oxidized fatty acids derived from arachidonate such as (15S)-HETE, eicosapentaenoate (EPA) such as (18R)- and (18S)-HEPE or docosahexaenoate (DHA) which lead to the formation of specialized pro-resolving mediators (SPM) lipoxin and resolvins E and D respectively, therefore it participates in anti-inflammatory responses (PubMed:<a href="http://www.uniprot.org/citations/17114001" target="\_blank">17114001</a>, PubMed:<a href="http://www.uniprot.org/citations/21206090" target="\_blank">21206090</a>, PubMed:<a href="http://www.uniprot.org/citations/31664810" target="\_blank">31664810</a>, PubMed:<a href="http://www.uniprot.org/citations/32404334" target="\_blank">32404334</a>, PubMed:<a href="http://www.uniprot.org/citations/8615788" target="\_blank">8615788</a>). Oxidation of DHA directly inhibits endothelial cell proliferation and sprouting angiogenesis via peroxisome proliferator-activated receptor gamma (PPARgamma) (By similarity). It does not catalyze the oxygenation of linoleic acid and does not convert (5S)-HETE to lipoxin isomers (PubMed:<a href="http://www.uniprot.org/citations/31664810" target="\_blank">31664810</a>). In addition to inflammatory processes, it participates in dendritic cell migration, wound healing through an antioxidant mechanism based on heme oxygenase-1 (HO-1) regulation expression, monocyte adhesion to the endothelium via ITGAM expression on monocytes (By similarity). Moreover, it helps establish an adaptive humoral

immunity by regulating primary resting B cells and follicular helper T cells and participates in the CD40-induced production of reactive oxygen species (ROS) after CD40 ligation in B cells through interaction with PIK3R1 that bridges ALOX5 with CD40 (PubMed:<a href="http://www.uniprot.org/citations/21200133" target="\_blank">21200133</a>). May also play a role in glucose homeostasis, regulation of insulin secretion and palmitic acid-induced insulin resistance via AMPK (By similarity). Can regulate bone mineralization and fat cell differentiation increases in induced pluripotent stem cells (By similarity).

#### Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:P48999, ECO:0000269|PubMed:18978352}. Nucleus matrix. Nucleus membrane; Peripheral membrane protein. Cytoplasm, perinuclear region. Cytoplasm, cytosol. Nucleus envelope. Nucleus intermembrane space. Note=Shuttles between cytoplasm and nucleus (PubMed:19233132). Found exclusively in the nucleus, when phosphorylated on Ser-272 (PubMed:18978352). Calcium binding promotes translocation from the cytosol and the nuclear matrix to the nuclear envelope and membrane association (PubMed:16275640, PubMed:19233132, PubMed:3118366, PubMed:8245774).

#### Anti-ALOX5 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-ALOX5 Antibody - Images



Anti-ALOX5 antibody, ABO10959, Western blotting All lanes: Anti ALOX5 (ABO10959) at 0.5ug/ml Lane 1: SW620 Whole Cell Lysate at 40ug Lane 2: JURKAT Whole Cell Lysate at 40ug Lane 3: COLO320 Whole Cell Lysate at 40ug Lane 4: A549 Whole Cell Lysate at 40ug Lane 5: MCF-7 Whole Cell Lysate at 40ug Predicted bind size: 78KD Observed bind size: 78KD



### **Anti-ALOX5 Antibody - Background**

ALOX5 (ARACHIDONATE 5-LIPOXYGENASE), also known as LOG5 or 5-LO (5-LIPOXYGENASE), is an enzyme that in humans is encoded by the ALOX5 gene. ALOX5 is a member of the lipoxygenase family of enzymes which also transforms EFAs into leukotrienes and is a current target for pharmaceutical intervention in a number of diseases. The enzyme 5-lipoxygenase catalyzes 2 reactions in the formation of leukotrienes. The ALOX5 gene is mapped to chromosome 10q11.21 based on an alignment of the ALOX5 sequence with the genomic sequence. Human 5-LO contains 3 nuclear localization sequences (NLSs) and a phosphorylation site involved in nuclear localization. Compared with age-matched 5-LO competent mice, the 5-LO knockout mice developed less right heart hypertrophy. Pharmacologic inhibition or ALOX5 gene disruption resulted in a significant decrease of beta-amyloid production and gamma-secretase levels. ALOX5 activity is short-lived, apparently in part because of an intrinsic instability of the enzyme.