

**Anti-IL-4 Antibody**  
Catalog # ABO11058**Specification**

---

**Anti-IL-4 Antibody - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">P05112</a>
Host	<b>Rabbit</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Format	<b>Lyophilized</b>

**Description**

Rabbit IgG polyclonal antibody for Interleukin-4(IL4) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-IL-4 Antibody - Additional Information**

**Gene ID** 3565

**Other Names**

Interleukin-4, IL-4, B-cell stimulatory factor 1, BSF-1, Binetrakin, Lymphocyte stimulatory factor 1, Pitrakinra, IL4

**Calculated MW**

17492 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br><br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted.

**Protein Name**

Interleukin-4(IL-4)

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human IL-4(72-88aa, AATVLRQFYSHHEKDTR).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

#### Sequence Similarities

Belongs to the IL-4/IL-13 family.

### Anti-IL-4 Antibody - Protein Information

Name IL4

#### Function

Cytokine secreted primarily by mast cells, T-cells, eosinophils, and basophils that plays a role in regulating antibody production, hematopoiesis and inflammation, and the development of effector T-cell responses (PubMed: [1993171](http://www.uniprot.org/citations/1993171), PubMed: [3016727](http://www.uniprot.org/citations/3016727)). Induces the expression of class II MHC molecules on resting B-cells. Enhances both secretion and cell surface expression of IgE and IgG1 (PubMed: [1993171](http://www.uniprot.org/citations/1993171)). Regulates also the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes (PubMed: [2521231](http://www.uniprot.org/citations/2521231)). Positively regulates IL31RA expression in macrophages. Stimulates autophagy in dendritic cells by interfering with mTORC1 signaling and through the induction of RUFY4. In addition, plays a critical role in higher functions of the normal brain, such as memory and learning (By similarity). Upon binding to IL4, IL4R receptor dimerizes either with the common IL2R gamma chain/IL2RG to produce the type 1 signaling complex, located mainly on hematopoietic cells, or with the IL13RA1 to produce the type 2 complex, which is expressed also on nonhematopoietic cells (PubMed: [10219247](http://www.uniprot.org/citations/10219247), PubMed: [11526337](http://www.uniprot.org/citations/11526337), PubMed: [18243101](http://www.uniprot.org/citations/18243101)). Engagement of both types of receptors initiates JAK3 and to a lower extent JAK1 phosphorylation leading to activation of the signal transducer and activator of transcription 6/STAT6 (PubMed: [7721895](http://www.uniprot.org/citations/7721895)).

#### Cellular Location


Secreted.

### Anti-IL-4 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-IL-4 Antibody - Images



Anti-IL-4 antibody, ABO11058, Western blotting All lanes: Anti IL-4 (ABO11058) at 0.5ug/ml  
 Lane 1: Recombinant Human IL-4 Protein 10ng  
 Lane 2: Recombinant Human IL-4 Protein 5ng  
 Lane 3: Recombinant Human IL-4 Protein 2.5ng  
 Predicted bind size: 14KD  
 Observed bind size: 14KD



Anti-IL-4 antibody, ABO11058, IHC(P) IHC(P): Human Tonsil Tissue

### Anti-IL-4 Antibody - Background

Interleukin-4(IL-4), also known as a B-cell stimulatory factor 1(BSF1), is an immunomodulatory cytokine, which can inhibit the growth of tumour cells. The human cDNA contains a single open reading frame encoding a protein of 153 amino acids, including a putative signal peptide. IL-4 may act as an autocrine growth factor in pancreatic cancer cells and also give rise to the possibility that cancer-derived IL-4 may suppress cancer-directed immunosurveillance in vivo in addition to its growth-promoting effects, thereby facilitating pancreatic tumor growth and metastasis. The mouse and human genes and their protein products show structural and functional similarities. The human IL-4 gene, which occurs as a single copy in the haploid genome, is mapped on chromosome 5. The standard product used in this kit is recombinant human IL-4, consisting of 130 amino acids with the molecular mass of 14KDa.