

**Anti-TrkC Antibody**  
Catalog # ABO11300**Specification**

---

**Anti-TrkC Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q16288</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for NT-3 growth factor receptor(NTRK3) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-TrkC Antibody - Additional Information**

**Gene ID** 4916

**Other Names**

NT-3 growth factor receptor, 2.7.10.1, GP145-TrkC, Trk-C, Neurotrophic tyrosine kinase receptor type 3, TrkC tyrosine kinase, NTRK3, TRKC

**Calculated MW**

94428 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat<br>

**Subcellular Localization**

Membrane; Single-pass type I membrane protein.

**Tissue Specificity**

Widely expressed but mainly in nervous tissue. Isoform 2 is expressed at higher levels in adult brain than in fetal brain.

**Protein Name**

NT-3 growth factor receptor

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human TrkC(172-186aa QLWQEQGEAKLNSQN), different from the related rat and mouse sequences by three amino acids.

### Purification

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### Sequence Similarities

Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.

## Anti-TrkC Antibody - Protein Information

**Name** NTRK3

**Synonyms** TRKC

### Function

Receptor tyrosine kinase involved in nervous system and probably heart development. Upon binding of its ligand NTF3/neurotrophin-3, NTRK3 autophosphorylates and activates different signaling pathways, including the phosphatidylinositol 3-kinase/AKT and the MAPK pathways, that control cell survival and differentiation.

### Cellular Location

Membrane; Single-pass type I membrane protein.

### Tissue Location


Widely expressed but mainly in nervous tissue. Isoform 2 is expressed at higher levels in adult brain than in fetal brain


## Anti-TrkC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-TrkC Antibody - Images





Anti-TrkC antibody, ABO11300, Western blotting Lane 1: Rat Brain Tissue Lysate Lane 2: Mouse Brain Tissue Lysate Lane 3: U87 Cell Lysate

### Anti-TrkC Antibody - Background

NTRK3 (Neurotrophic Tyrosine Kinase Receptor Type 3), also known as TRKC, is a protein that in humans is encoded by the NTRK3 gene. By PCR analysis of a somatic cell hybrid panel and by fluorescence in situ hybridization with the cDNA clone, McGregor et al. (1994) mapped the NTRK3 gene to 15q24-q25. Lamballe et al. (1991) isolated and characterized TRKC, a member of the TRK family of tyrosine protein kinase genes. They found that TRKC is preferentially expressed in the brain; in situ hybridization studies showed transcripts in the hippocampus, cerebral cortex, and the granular cell layer of the cerebellum. By functional studies in HeLa cells, Muinos-Gimeno et al. (2009) demonstrated that 5 miRNAs regulate the truncated form of NTRK3.