

Anti-TRAF2 Antibody
Catalog # ABO11314

Specification

Anti-TRAF2 Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	Q12933
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for TNF receptor-associated factor 2 (TRAF2) detection. Tested with WB, IHC-P, ICC in Human; Mouse; Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-TRAF2 Antibody - Additional Information

Gene ID 7186

Other Names

TNF receptor-associated factor 2, 6.3.2.-, E3 ubiquitin-protein ligase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2, TRAP3

Calculated MW

55859 MW KDa

Application Details

Immunocytochemistry , 0.5-1 µg/ml, Human, -
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Cytoplasm .

Protein Name

TNF receptor-associated factor 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human TRAF2(305-325aa RQHRLDQDKIEALSSKVQLE), different from the related rat and mouse sequences by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r^oConstitution, at 4°C for one month. It^oCan also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the TNF receptor-associated factor family. A subfamily.

Anti-TRAF2 Antibody - Protein Information

Name TRAF2 {ECO:0000303|PubMed:28489822, ECO:0000312|HGNC:HGNC:12032}

Function

E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:10346818, PubMed:11784851, PubMed:12917689, PubMed:15383523, PubMed:18981220, PubMed:19150425, PubMed:19810754, PubMed:19918265, PubMed:19937093, PubMed:20047764, PubMed:20064526, PubMed:20385093, PubMed:20577214, PubMed:22212761). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:23453969, PubMed:28489822). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:15383523, PubMed:18981220). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:11907583, PubMed:19506082). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:15121867). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:23453969). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:28489822). Required for normal antibody isotype switching from IgM to IgG (By similarity).


Cellular Location
Cytoplasm

Anti-TRAF2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-TRAF2 Antibody - Images



Anti-TRAF2 antibody, ABO11314, Western blotting
Recombinant Protein Detection Source: E.coli derived -recombinant Human TRAF2, 30.4KD(162aa tag+ Q261-V369)
Lane 1: Recombinant Human TRAF2 Protein 10ng
Lane 2: Recombinant Human TRAF2 Protein 5ng
Lane 3: Recombinant Human TRAF2 Protein 2.5ng



Anti-TRAF2 antibody, ABO11314, IHC(P)IHC(P): Human Tonsil Tissue



Anti-TRAF2 antibody, ABO11314, ICCICC: HELA Cell

Anti-TRAF2 Antibody - Background

TRAF2(TNF Receptor-Associated Factor 2), also called TRAP, is a protein that in humans is encoded by the TRAF2 gene. The protein encoded by this gene is a member of the TNF receptor(TNFR) associated factor(TRAF) protein family. TRAF2 is a common signal transducer for TNFR2 and CD40 that mediates activation of NF-kappa-B. Rothe et al.(1996) identified ITRAF, which binds to TRAF1, TRAF2, and TRAF3, and that when overexpressed inhibits TRAF2-mediated NF-kappa-B activation. They proposed that ITRAF is an inhibitor of TRAF function that regulates TRAF protein activity by sequestering TRAFs in a latent state in the cytoplasm. Kanamori et al.(2002) found that mouse Traf2 interacted directly with T2bp, and they presented evidence that T2BP is involved in TNF-mediated signaling by its interaction with TRAF2.