

Anti-PLK2 Antibody
Catalog # ABO11439**Specification****Anti-PLK2 Antibody - Product Information**

Application	WB, IHC
Primary Accession	O9NYY3
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Serine/threonine-protein kinase PLK2(PLK2) detection. Tested with WB, IHC-P, ICC in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-PLK2 Antibody - Additional Information

Gene ID 10769

Other Names

Serine/threonine-protein kinase PLK2, 2.7.11.21, Polo-like kinase 2, PLK-2, hPlk2, Serine/threonine-protein kinase SNK, hSNK, Serum-inducible kinase, PLK2, SNK

Calculated MW

Serine/threonine-protein kinase PLK2 KDa

Application Details

Immunocytochemistry , 0.5-1 µg/ml, Human, Mouse, Rat
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Tumor suppressor serine/threonine-protein kinase involved in synaptic plasticity, centriole duplication and G1/S phase transition. Polo-like kinases act by binding and phosphorylating proteins that are already phosphorylated on a specific motif recognized by the POLO box domains. Phosphorylates CENPJ, NPM1, RAPGEF2, RASGRF1, SNCA, SIPA1L1 and SYNGAP1. Plays a key role in synaptic plasticity and memory by regulating the Ras and Rap protein signaling: required for overactivity-dependent spine remodeling by phosphorylating the Ras activator RASGRF1 and the Rap inhibitor SIPA1L1 leading to their degradation by the proteasome. Conversely, phosphorylates the Rap activator RAPGEF2 and the Ras inhibitor SYNGAP1, promoting their activity. Also regulates synaptic plasticity independently of kinase activity, via its interaction with NSF that disrupts the interaction between NSF and the GRIA2 subunit of AMPARs, leading to a rapid rundown of AMPAR-mediated current that occludes long term depression. Required for procentriole formation and centriole duplication by phosphorylating CENPJ and NPM1, respectively. Its induction by p53/TP53 suggests that it may participate in the mitotic checkpoint following stress. .

Tissue Specificity

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole . Cell projection, dendrite . Localizes to centrosomes during early G1 phase where it only associates to the mother centriole and then distributes equally to both mother and daughter centrioles at the onset of S phase.

Source

Expressed at higher level in the fetal lung, kidney, spleen and heart. .

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human PLK2(666-685aa SSELKNRMEYALNMLLQRCN), different from the related mouse sequence by one amino acid, and from the related rat sequence by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.

Anti-PLK2 Antibody - Protein Information

Name PLK2

Synonyms SNK

Function

Tumor suppressor serine/threonine-protein kinase involved in synaptic plasticity, centriole duplication and G1/S phase transition. Polo-like kinases act by binding and phosphorylating proteins that are already phosphorylated on a specific motif recognized by the POLO box domains. Phosphorylates CENPJ, NPM1, RAPGEF2, RASGRF1, SNCA, SIPA1L1 and SYNGAP1. Plays a key role in synaptic plasticity and memory by regulating the Ras and Rap protein signaling: required for overactivity-dependent spine remodeling by phosphorylating the Ras activator RASGRF1 and the Rap inhibitor SIPA1L1 leading to their degradation by the proteasome. Conversely, phosphorylates the Rap activator RAPGEF2 and the Ras inhibitor SYNGAP1, promoting their activity. Also regulates synaptic plasticity independently of kinase activity, via its interaction with NSF that disrupts the interaction between NSF and the GRIA2 subunit of AMPARs, leading to a rapid rundown of AMPAR-mediated current that occludes long term depression. Required for procentriole formation and centriole duplication by phosphorylating CENPJ and NPM1, respectively. Its induction by p53/TP53 suggests that it may participate in the mitotic checkpoint following stress.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cell projection,

dendrite Note=Localizes to centrosomes during early G1 phase where it only associates to the mother centriole and then distributes equally to both mother and daughter centrioles at the onset of S phase

Tissue Location


Expressed at higher level in the fetal lung, kidney, spleen and heart.

Anti-PLK2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-PLK2 Antibody - Images




Anti-PLK2 antibody, ABO11439, Western blotting Lane 1: A431 Cell Lysate Lane 2: 293T Cell Lysate Lane 3: COLO320 Cell Lysate



Anti-PLK2 antibody, ABO11439, IHC(P)IHC(P): Human Lung Cancer Tissue



Anti-PLK2 antibody, ABO11439, IHC(P)IHC(P): Rat Intestine Tissue



Anti-PLK2 antibody, ABO11439, IHC(P)IHC(P): Rat Brain Tissue

Anti-PLK2 Antibody - Background

Polo-like kinase2, also known as SNK is an enzyme that in humans is encoded by the PLK2 gene. PLK2 is a member of the polo" family of serine/threonine protein kinases that have a role in normal cell division. The International Radiation Hybrid Mapping Consortium mapped the SNK gene to chromosome 5. SNK involved in synaptic plasticity