

**Anti-FABP6 Antibody**  
Catalog # ABO11469**Specification****Anti-FABP6 Antibody - Product Information**

Application	IHC, WB
Primary Accession	<a href="#">P51161</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Gastrotropin(FABP6) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-FABP6 Antibody - Additional Information**

**Gene ID** 2172

**Other Names**

Gastrotropin, GT, Fatty acid-binding protein 6, Ileal lipid-binding protein, ILBP, Intestinal 15 kDa protein, I-15P, Intestinal bile acid-binding protein, I-BABP, FABP6, ILBP, ILLBP

**Calculated MW**

14371 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Isoform 1: Cytoplasm.

**Tissue Specificity**

Isoform 2 is expressed in colorectal adenocarcinomas and their adjacent normal mucosa (at protein level). Isoform 1 is expressed in the jejunum, ileum, cecum and ascending colon intestine. Isoform 2 is expressed in the gallbladder, duodenum, jejunum, ileum, cecum, ascending, transverse and descending colon, sigmoid colon and rectum. .

**Protein Name**

Gastrotropin

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human FABP6(15-33aa YDEFMKLLGISSDVIEKAR).

#### **Purification**

Immunogen affinity purified.

#### **Cross Reactivity**

No cross reactivity with other proteins

#### **Storage**

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

#### **Sequence Similarities**

Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.

### **Anti-FABP6 Antibody - Protein Information**

**Name** FABP6

**Synonyms** ILBP, ILLBP

#### **Function**

Binds to bile acids and is involved in enterohepatic bile acid metabolism. Required for efficient apical to basolateral transport of conjugated bile acids in ileal enterocytes (By similarity). In vitro binds to bile acids in the order: deoxycholic acid > cholic acid > chenodeoxycholic acid and respective BA conjugation modifies affinities in the order taurine-conjugated > glycine-conjugated > unconjugated bile acids. Stimulates gastric acid and pepsinogen secretion (By similarity).

#### **Cellular Location**

[Isoform 1]: Cytoplasm {ECO:0000250|UniProtKB:P80020}. Membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P50119}; Cytoplasmic side {ECO:0000250|UniProtKB:P50119}

#### **Tissue Location**


Isoform 1 is expressed in the jejunum, ileum, cecum and ascending colon intestine. Isoform 2 is xpressed in the gallbladder, duodenum, jejunum, ileum, cecum, ascending, transverse and descending colon, sigmoid colon and rectum. Isoform 2 is expressed in colorectal adenocarcinomas and their adjacent normal mucosa (at protein level).

### **Anti-FABP6 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-FABP6 Antibody - Images**



Anti-FABP6 antibody, ABO11469, IHC(P)IHC(P): Human Lung Cancer Tissue



Anti-FABP6 antibody, ABO11469, Western blottingAll lanes: Anti FABP6 (ABO11469) at 0.5ug/mlLane 1: COLO320 Whole Cell Lysate at 40ugLane 2: SW620 Whole Cell Lysate at 40ugPredicted bind size: 14KDObserved bind size: 14KD

**Anti-FABP6 Antibody - Background**

Fatty acid binding protein 6, ileal, also called FABP6 or ILBP is a protein which in humans is encoded by the FABP6 gene. Ileal lipid-binding protein(ILBP) is a member of a family of intracellular fatty acid, retinoid, and bile acid-binding proteins. This gene is mapped to 5q33.3. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. FABP6 and FABP1(the liver fatty acid binding protein) are also able to bind bile acids. It is though that FABPs roles include fatty acid uptake, transport, and metabolism.