

**Anti-UBE2I/UBC9 Antibody**  
Catalog # ABO11568**Specification****Anti-UBE2I/UBC9 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P63279</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for SUMO-conjugating enzyme UBC9(UBE2I) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-UBE2I/UBC9 Antibody - Additional Information**

**Gene ID** 7329

**Other Names**

SUMO-conjugating enzyme UBC9, 2.3.2.-, RING-type E3 SUMO transferase UBC9, SUMO-protein ligase, Ubiquitin carrier protein 9, Ubiquitin carrier protein I, Ubiquitin-conjugating enzyme E2 I, Ubiquitin-protein ligase I, p18, UBE2I, UBC9, UBCE9

**Calculated MW**

18007 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Rat, Human, Mouse

**Subcellular Localization**

Nucleus. Cytoplasm. Mainly nuclear. In spermatocytes, localizes in synaptonemal complexes. Recruited by BCL11A into the nuclear body (By similarity).

**Tissue Specificity**

Expressed in heart, skeletal muscle, pancreas, kidney, liver, lung, placenta and brain. Also expressed in testis and thymus.

**Protein Name**

SUMO-conjugating enzyme UBC9

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human UBE2I(108-122aa

IKQILLGIQELLNE), identical to the related mouse and rat sequences.

#### **Purification**

Immunogen affinity purified.

#### **Cross Reactivity**

No cross reactivity with other proteins

#### **Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

#### **Sequence Similarities**

Belongs to the ubiquitin-conjugating enzyme family.

### **Anti-UBE2I/UBC9 Antibody - Protein Information**

**Name** UBE2I

**Synonyms** UBC9, UBCE9

#### **Function**

Accepts the ubiquitin-like proteins SUMO1, SUMO2, SUMO3, SUMO4 and SUMO1P1/SUMO5 from the UBLE1A-UBLE1B E1 complex and catalyzes their covalent attachment to other proteins with the help of an E3 ligase such as RANBP2, CBX4 and ZNF451. Can catalyze the formation of poly-SUMO chains. Necessary for sumoylation of FOXL2 and KAT5. Essential for nuclear architecture and chromosome segregation. Sumoylates p53/TP53 at 'Lys-386'. Mediates sumoylation of ERCC6 which is essential for its transcription-coupled nucleotide excision repair activity (PubMed: [26620705](http://www.uniprot.org/citations/26620705)).

#### **Cellular Location**

Nucleus. Cytoplasm Cytoplasm, perinuclear region Note=Mainly nuclear (By similarity). In spermatocytes, localizes in synaptonemal complexes (PubMed:8610150). Recruited by BCL11A into the nuclear body (By similarity). {ECO:0000250|UniProtKB:P63280, ECO:0000269|PubMed:8610150}

#### **Tissue Location**

Expressed in heart, skeletal muscle, pancreas, kidney, liver, lung, placenta and brain. Also expressed in testis and thymus.


### **Anti-UBE2I/UBC9 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

### Anti-UBE2I/UBC9 Antibody - Images



Anti-UBE2I/UBC9 antibody, ABO11568, All Western blotting  
All lanes: Anti-UBE2I(ABO11568) at 0.5ug/ml  
Lane 1: Rat Cardiac Muscle Tissue Lysate at 40ug  
Lane 2: Rat Skeletal Muscle Tissue Lysate at 40ug  
Lane 3: Rat Pancreas Tissue Lysate at 40ug  
Predicted bind size: 18KD  
Observed bind size: 18KD

### Anti-UBE2I/UBC9 Antibody - Background

SUMO-conjugating enzyme UBC9(UBE2I), also called UBC9, is a protein that in humans is encoded by the UBE2I gene. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. It is mapped to 16p13.3. UBC9 could fully complement the mutant phenotype of a yeast *ubc9* mutant strain. This gene may play a similar role via interaction with WT1, which is able to impose a block to cell cycle progression in eukaryotic cells. What's more, it could support the growth of yeast *ubc9* temperature-sensitive mutants at nonpermissive temperatures, indicating that the gene is a functional homolog of yeast *ubc9*. UBC9 is specifically associated with FHIT, such as FHIT may be involved in cell cycle control through its interaction with UBC9.