

**Anti-Mucin 5AC Antibody**  
Catalog # ABO11611**Specification****Anti-Mucin 5AC Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P98088</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Mucin-5AC(MUC5AC) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Mucin 5AC Antibody - Additional Information**

Gene ID 4586

**Other Names**

Mucin-5AC, MUC-5AC, Gastric mucin, Lewis B blood group antigen, LeB, Major airway glycoprotein, Mucin-5 subtype AC, tracheobronchial, Tracheobronchial mucin, TBM, MUC5AC  
{ECO:0000303|PubMed:11535137, ECO:0000312|HGNC:HGNC:7515}

**Calculated MW**

585570 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted .

**Tissue Specificity**

Highly expressed in surface mucosal cells of respiratory tract and stomach epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal duodenum. .

**Protein Name**

Mucin-5AC

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human Mucin 5AC(552-567aa TMQLFMQLAPKLRGQT).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Contains 1 CTCK (C-terminal cystine knot-like) domain.

**Anti-Mucin 5AC Antibody - Protein Information**

**Name** MUC5AC {ECO:0000303|PubMed:11535137, ECO:0000312|HGNC:HGNC:7515}

**Function**

Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system (PubMed:<a href="http://www.uniprot.org/citations/14535999" target="\_blank">14535999</a>, PubMed:<a href="http://www.uniprot.org/citations/14718370" target="\_blank">14718370</a>). Interacts with H.pylori in the gastric epithelium, Barrett's esophagus as well as in gastric metaplasia of the duodenum (GMD) (PubMed:<a href="http://www.uniprot.org/citations/14535999" target="\_blank">14535999</a>).

**Cellular Location**

Secreted

**Tissue Location**


Highly expressed in surface mucosal cells of respiratory tract and stomach epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal duodenum.

**Anti-Mucin 5AC Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Mucin 5AC Antibody - Images**



Anti-Mucin 5AC antibody, ABO11611, All Western blotting  
All lanes: Anti-MUC5AC(ABO11611) at 0.5ug/ml  
Lane 1: SGC Whole Cell Lysate at 40ug  
Lane 2: HELA Whole Cell Lysate at 40ug  
Lane 3: COLO320 Whole Cell Lysate at 40ug  
Lane 4: SW620 Whole Cell Lysate at 40ug  
Lane 5: MCF-7 Whole Cell Lysate at 40ug  
Lane 6: PANC Whole Cell Lysate at 40ug  
Predicted bind size: 527KD  
Observed bind size: 300KD

#### **Anti-Mucin 5AC Antibody - Background**

Mucin-5AC is a protein that in humans is encoded by the MUC5AC gene. It is mapped to 11p15.5. MUC5AC is highly expressed in adult trachea and stomach. This gene has been linked to mucus hypersecretion in the pulmonary tracts and associated to chronic obstructive pulmonary disease (COPD). It has been shown that activation of the NF-kappa-B signaling pathway by inflammatory cytokine TNF-alpha upregulated MUC5AC mRNA expression nearly 6-fold. MUC5AC is also a direct and critical mediator of resistance during intestinal nematode infection.