

Anti-ADFP Picoband Antibody
Catalog # ABO11701**Specification**

Anti-ADFP Picoband Antibody - Product Information

Application	WB, IHC
Primary Accession	Q99541
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Perilipin-2 (PLIN2) detection. Tested with WB, IHC-P in Human; Mouse; Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-ADFP Picoband Antibody - Additional Information

Gene ID 123

Other Names

Perilipin-2, Adipophilin, Adipose differentiation-related protein, ADRP, PLIN2, ADFP

Calculated MW

48075 MW KDa

Application Details

Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Membrane; Peripheral membrane protein.

Tissue Specificity

Milk lipid globules.

Protein Name

Perilipin-2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

E. coli-derived human ADFP recombinant protein (Position: K226-Q418). Human ADFP shares 88.4% amino acid (aa) sequence identity with mouse ADFP.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-ADFP Picoband Antibody - Protein Information

Name PLIN2 ([HGNC:248](#))

Synonyms ADFP

Function

Structural component of lipid droplets, which is required for the formation and maintenance of lipid storage droplets.

Cellular Location

Membrane {ECO:0000250|UniProtKB:P43883}; Peripheral membrane protein {ECO:0000250|UniProtKB:P43883}. Lipid droplet


Tissue Location


Milk lipid globules..

Anti-ADFP Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-ADFP Picoband Antibody - Images




Western blot analysis of ADFP expression in MCF-7 whole cell lysates (lane 1). ADFP at 48KD was detected using rabbit anti- ADFP Antigen Affinity purified polyclonal antibody (Catalog # ABO11701) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



ADFP was detected in paraffin-embedded sections of mouse testis tissues using rabbit anti- ADFP Antigen Affinity purified polyclonal antibody (Catalog # ABO11701) at 1 1/4 µg/mL. The immunohistochemical section was developed using SABC method .



ADFP was detected in paraffin-embedded sections of rat testis tissues using rabbit anti- ADFP Antigen Affinity purified polyclonal antibody (Catalog # ABO11701) at 1 1/4 µg/mL. The immunohistochemical section was developed using SABC method .



ADFP was detected in paraffin-embedded sections of human mammary cancer tissues using rabbit anti- ADFP Antigen Affinity purified polyclonal antibody (Catalog # ABO11701) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Anti-ADFP Picoband Antibody - Background

Adipose differentiation-related protein, also known as perilipin 2 (PLIN2), ADRP or adipophilin, is a protein which in humans is encoded by the ADFP gene. The protein encoded by this gene belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This protein is associated with the lipid globule surface membrane material, and maybe involved in development and maintenance of adipose tissue. However, it is not restricted to adipocytes as previously thought, but is found in a wide range of cultured cell lines, including fibroblasts, endothelial and epithelial cells, and tissues, such as lactating mammary gland, adrenal cortex, Sertoli and Leydig cells, and hepatocytes in alcoholic liver cirrhosis, suggesting that it may serve as a marker of lipid accumulation in diverse cell types and diseases. Alternatively spliced transcript variants have been found for this gene.