

Anti-COX1/Cyclooxygenase 1 Picoband Antibody
Catalog # ABO11727

Specification

Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Product Information

Application	IHC
Primary Accession	P23219
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Prostaglandin G/H synthase 1 (PTGS1) detection. Tested with WB, IHC-P in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Additional Information

Gene ID 5742

Other Names

Prostaglandin G/H synthase 1, 1.14.99.1, Cyclooxygenase-1, COX-1, Prostaglandin H2 synthase 1, PGH synthase 1, PGHS-1, PHS 1, Prostaglandin-endoperoxide synthase 1, PTGS1, COX1

Calculated MW

68686 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein.

Protein Name

Prostaglandin G/H synthase 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

E.coli-derived human COX1 recombinant protein (Position: E318-L599). Human COX1 shares 90% and 89% amino acid (aa) sequences identity with mouse and rat COX1, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the prostaglandin G/H synthase family.

Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Protein Information

Name PTGS1 ([HGNC:9604](#))

Function

Dual cyclooxygenase and peroxidase that plays an important role in the biosynthesis pathway of prostanoids, a class of C20 oxylipins mainly derived from arachidonate ((5Z,8Z,11Z,14Z)-eicosatetraenoate, AA, C20:4(n-6)), with a particular role in the inflammatory response. The cyclooxygenase activity oxygenates AA to the hydroperoxy endoperoxide prostaglandin G2 (PGG2), and the peroxidase activity reduces PGG2 to the hydroxy endoperoxide prostaglandin H2 (PGH2), the precursor of all 2-series prostaglandins and thromboxanes. This complex transformation is initiated by abstraction of hydrogen at carbon 13 (with S-stereochemistry), followed by insertion of molecular O2 to form the endoperoxide bridge between carbon 9 and 11 that defines prostaglandins. The insertion of a second molecule of O2 (bis-oxygenase activity) yields a hydroperoxy group in PGG2 that is then reduced to PGH2 by two electrons (PubMed:7947975). Involved in the constitutive production of prostanoids in particular in the stomach and platelets. In gastric epithelial cells, it is a key step in the generation of prostaglandins, such as prostaglandin E2 (PGE2), which plays an important role in cytoprotection. In platelets, it is involved in the generation of thromboxane A2 (TXA2), which promotes platelet activation and aggregation, vasoconstriction and proliferation of vascular smooth muscle cells (Probable). Can also use linoleate (LA, (9Z,12Z)- octadecadienoate, C18:2(n-6)) as substrate and produce hydroxyoctadecadienoates (HODEs) in a regio- and stereospecific manner, being (9R)-HODE ((9R)-hydroxy-(10E,12Z)-octadecadienoate) and (13S)- HODE ((13S)-hydroxy-(9Z,11E)-octadecadienoate) its major products (By similarity).

Cellular Location


Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein

Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Images




Anti-COX1 Picoband antibody, ABO11727-1.JPGIHC(P): Human Intestinal Cancer Tissue




Anti-COX1 Picoband antibody, ABO11727-2.JPGIHC(P): Human Lung Cancer Tissue



Anti-COX1 Picoband antibody, ABO11727-3.JPGIHC(P): Rat Intestine Tissue



Anti-COX1 Picoband antibody, ABO11727-4.JPGIHC(P): Human Mammary Cancer Tissue



Anti-COX1 Picoband antibody, ABO11727-5.jpg All lanes: Anti-COX1 (ABO11727) at 0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 40ug
Lane 2: HELA Whole Cell Lysate at 40ug
Lane 3: JURKAT Whole Cell Lysate at 40ug
Lane 4: K562 Whole Cell Lysate at 40ug
Lane 5: MCF-7 Whole Cell Lysate at 40ug
Lane 6: A549 Whole Cell Lysate at 40ug
Predicted bind size: 69KD
Observed bind size: 69KD

Anti-COX1/Cyclooxygenase 1 Picoband Antibody - Background

Cyclooxygenase 1 (COX1), also known as Prostaglandin-endoperoxide synthase (PTGS1) or mitochondrial cytochrome c oxidase subunit 1, is the key enzyme in prostaglandin biosynthesis. The gene was approximately 40 kb long, with 11 protein-coding exons. There were 599 amino acid residues with a calculated molecular mass of approximately 68 kD. By analysis of a human/hamster somatic hybrid DNA panel, it was demonstrated that the PTGS1 gene maps to chromosome 9. Human prostaglandin endoperoxide synthase exhibited 91% amino acid identity with the sheep enzyme. Prostaglandin synthase 1 gene disruption in mice reduces arachidonic acid-induced inflammation and indomethacin-induced gastric ulceration.