

Anti-CD23/FCER2 Antibody
Catalog # ABO11772

Specification

Anti-CD23/FCER2 Antibody - Product Information

Application	IHC, WB
Primary Accession	P20693
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Low affinity immunoglobulin epsilon Fc receptor(FCER2) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CD23/FCER2 Antibody - Additional Information

Gene ID 14128

Other Names

Low affinity immunoglobulin epsilon Fc receptor, Fc-epsilon-RII, Lymphocyte IgE receptor, CD23, Fcer2, Fcer2a

Calculated MW

Langerhans cells KDa

Application Details

Immunohistochemistry(Frozen Section), 0.5-1 µg/ml, Mouse,
-
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By
Heat
Western blot, 0.1-0.5 µg/ml, Mouse

Subcellular Localization

and macrophages. As part of a mapping of multiple probes to specific bands on chromosome 19 by fluorescence in situ hybridization

Tissue Specificity

the FCE2 gene was assigned to 19p13.3. CD23 (FCE2) is a key molecule for B-cell activation and growth. It is the low-affinity receptor for IgE. The truncated molecule can be secreted

Source

then functioning as a potent mitogenic growth factor."

Protein Name

sc 13940|sc 31574|sc 7426|sc 365524

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived mouse CD23 recombinant protein (Position: E50-P331). Mouse CD23 shares 52% amino acid (aa) sequence identity with human CD23.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Low affinity immunoglobulin epsilon Fc receptor;Fc-epsilon-RII;Lymphocyte IgE receptor;CD23;Fcer2;Fcer2a;

Anti-CD23/FCER2 Antibody - Protein Information

Name Fcer2

Synonyms Fcer2a

Function

Low-affinity receptor for immunoglobulin E (IgE) and CR2/CD21. Has essential roles in the regulation of IgE production and in the differentiation of B cells. On B cells, initiates IgE-dependent antigen uptake and presentation to T cells. On macrophages, upon IgE binding and antigen cross-linking induces intracellular killing of parasites through activation of L-Arginine-nitric oxide pathway.


Cellular Location


Cell membrane; Single-pass type II membrane protein. Cell membrane; Lipid- anchor. Secreted {ECO:0000250|UniProtKB:P06734}

Anti-CD23/FCER2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-CD23/FCER2 Antibody - Images



Anti-CD23 Picoband antibody, ABO11772-1.JPGIHC(P): Human Tonsil Tissue



Anti-CD23 Picoband antibody, ABO11772-2.JPGIHC(P): Mouse Spleen Tissue



Anti-CD23 Picoband antibody, ABO11772-3.JPGIHC(P): Rat Spleen Tissue



Anti-CD23 Picoband antibody, ABO11772-4.jpg All lanes: Anti-CD23(ABO11772) at 0.5ug/ml WB:
Mouse Liver Tissue Lysate at 40ug Predicted bind size: 37KD Observed bind size: 37KD

Anti-CD23/FCER2 Antibody - Background

CD23, also known as Fc epsilon RII, or Fc ϵ RII, is the low-affinity" receptor for IgE