

Anti-GLUT4 Picoband Antibody
Catalog # ABO11815**Specification****Anti-GLUT4 Picoband Antibody - Product Information**

Application	IHC
Primary Accession	P14672
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Solute carrier family 2, facilitated glucose transporter member 4 (SLC2A4) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-GLUT4 Picoband Antibody - Additional Information

Gene ID 6517

Other Names

Solute carrier family 2, facilitated glucose transporter member 4, Glucose transporter type 4, insulin-responsive, GLUT-4, SLC2A4, GLUT4

Calculated MW

54787 MW KDa

Application Details

Immunohistochemistry(Frozen Section), 0.5-1 µg/ml, Mouse, Rat,
-
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By
Heat
Western blot, 0.1-0.5 µg/ml, Rat, Human

Subcellular Localization

Cell membrane ; Multi-pass membrane protein . Endomembrane system ; Multi-pass membrane protein . Cytoplasm, perinuclear region . Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized. The dileucine internalization motif is critical for intracellular sequestration.

Tissue Specificity

Skeletal and cardiac muscles; brown and white fat.

Protein Name

Solute carrier family 2, facilitated glucose transporter member 4

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E.coli-derived human GLUT4 recombinant protein (Position: N333-D509). Human GLUT4 shares 97% amino acid (aa) sequence identity with mouse GLUT4.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily.

Anti-GLUT4 Picoband Antibody - Protein Information

Name SLC2A4 ([HGNC:11009](#))

Function

Insulin-regulated facilitative glucose transporter, which plays a key role in removal of glucose from circulation. Response to insulin is regulated by its intracellular localization: in the absence of insulin, it is efficiently retained intracellularly within storage compartments in muscle and fat cells. Upon insulin stimulation, translocates from these compartments to the cell surface where it transports glucose from the extracellular milieu into the cell.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P14142}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P14142} Endomembrane system; Multi-pass membrane protein. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P14142}. Note=Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized (PubMed:8300557). The dileucine internalization motif is critical for intracellular sequestration (PubMed:8300557). Insulin stimulation induces translocation to the cell membrane (By similarity) {ECO:0000250|UniProtKB:P14142, ECO:0000269|PubMed:8300557}

Tissue Location

Skeletal and cardiac muscles; brown and white fat.


Anti-GLUT4 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)


Anti-GLUT4 Picoband Antibody - Images




Anti-GLUT4 Picoband antibody, ABO11815-1.jpg All lanes: Anti GLUT4 (ABO11815) at 0.5ug/ml
Lane 1: Rat Cardiac Muscle Tissue Lysate at 50ug
Lane 2: Rat Skeletal Muscle Tissue Lysate at 50ug
Predicted bind size: 55KD
Observed bind size: 55KD




Anti-GLUT4 Picoband antibody, ABO11815-2.JPG IHC(F): Rat Cardiac Muscle Tissue




Anti-GLUT4 Picoband antibody, ABO11815-3.JPGIHC(F): Mouse Cardiac Muscle Tissue



Anti-GLUT4 Picoband antibody, ABO11815-4.JPGIHC(P): Human Intestinal Cancer Tissue



Anti-GLUT4 Picoband antibody, ABO11815-5.JPGIHC(P): Rat Cardiac Muscle Tissue



Anti-GLUT4 Picoband antibody, ABO11815-6.JPGIHC(P): Mouse Skeletal Muscle Tissue

Anti-GLUT4 Picoband Antibody - Background

GLUT4, also known as SLC2A4 or solute carrier family 2 (facilitated glucose transporter) member 4, is a protein that in humans is encoded by the GLUT4 gene. It is mapped to 17p13.1. This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM).