

Anti-DHFR Picoband Antibody
Catalog # ABO11868**Specification****Anti-DHFR Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	P00374
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Dihydrofolate reductase(DHFR) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-DHFR Picoband Antibody - Additional Information

Gene ID 1719

Other Names

Dihydrofolate reductase, 1.5.1.3, DHFR

Calculated MW

21453 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Tissue Specificity

Widely expressed in fetal and adult tissues, including throughout the fetal and adult brains and whole blood. Expression is higher in the adult brain than in the fetal brain. .

Protein Name

Dihydrofolate reductase

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

E.coli-derived human DHFR recombinant protein (Position: V2-D187). Human DHFR shares 90% amino acid (aa) sequence identity with both mouse and rat DHFR.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the dihydrofolate reductase family.

Anti-DHFR Picoband Antibody - Protein Information

Name DHFR

Function

Key enzyme in folate metabolism. Contributes to the de novo mitochondrial thymidylate biosynthesis pathway. Catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis. Binds its own mRNA and that of DHFR2.

Cellular Location

Mitochondrion {ECO:0000250|UniProtKB:P00375}. Cytoplasm {ECO:0000250|UniProtKB:P00375}

Tissue Location


Widely expressed in fetal and adult tissues, including throughout the fetal and adult brains and whole blood Expression is higher in the adult brain than in the fetal brain


Anti-DHFR Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-DHFR Picoband Antibody - Images







Anti- DHFR Picoband antibody, ABO11868, Western blotting All lanes: Anti DHFR (ABO11868) at 0.5ug/ml WB: Recombinant Human DHFR Protein 0.5ng Predicted bind size: 32KD Observed bind size: 32KD




Anti- DHFR Picoband antibody, ABO11868, Western blotting All lanes: Anti DHFR (ABO11868) at 0.5ug/ml Lane 1: Rat Liver Tissue Lysate at 50ug Lane 2: Mouse Liver Tissue Lysate at 50ug Lane 3: HepG2 Whole Cell Lysate at 40ug Lane 4: Jurkat Whole Cell Lysate at 40ug Predicted bind size: 21KD Observed bind size: 21KD



Anti- DHFR Picoband antibody, ABO11868, IHC(P) IHC(P): Mouse Brain Tissue



Anti- DHFR Picoband antibody, ABO11868,IHC(P)IHC(P): Rat Brain Tissue



Anti- DHFR Picoband antibody, ABO11868,IHC(P)IHC(P): Human Lung Cancer Tissue

Anti-DHFR Picoband Antibody - Background

Dihydrofolate reductase, or DHFR, is an enzyme that reduces dihydrofolic acid to tetrahydrofolic acid, using NADPH as electron donor, which can be converted to the kinds of tetrahydrofolate cofactors used in 1-carbon transfer chemistry. In humans, the DHFR enzyme is encoded by the DHFR gene. It is found in the q11â†'q22 region of chromosome 5. What's more, DHFR belongs to the dihydrofolate reductase family, and it converts dihydrofolate into tetrahydrofolate, a methyl group shuttle required for the de novo synthesis of purines, thymidylic acid, and certain amino acids. DHFR is the key enzyme in folate metabolism. In addition, DHFR catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis.