

**Anti-Caspase-9 Picoband Antibody**  
Catalog # ABO12023**Specification****Anti-Caspase-9 Picoband Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P55211</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Caspase-9(CASP9) detection. Tested with WB in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Caspase-9 Picoband Antibody - Additional Information**

**Gene ID** 842

**Other Names**

Caspase-9, CASP-9, 3.4.22.62, Apoptotic protease Mch-6, Apoptotic protease-activating factor 3, APAF-3, ICE-like apoptotic protease 6, ICE-LAP6, Caspase-9 subunit p35, Caspase-9 subunit p10, CASP9, MCH6

**Calculated MW**

46281 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Tissue Specificity**

Ubiquitous, with highest expression in the heart, moderate expression in liver, skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart, specifically expressed in myocytes.

**Protein Name**

Caspase-9

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

E.coli-derived human Caspase-9 recombinant protein (Position: E3-D228). Human Caspase-9 shares 63% amino acid (aa) sequence identity with mouse Caspase-9.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

### Sequence Similarities

Belongs to the peptidase C14A family.

## Anti-Caspase-9 Picoband Antibody - Protein Information

**Name** CASP9

**Synonyms** MCH6

### Function

Involved in the activation cascade of caspases responsible for apoptosis execution. Binding of caspase-9 to Apaf-1 leads to activation of the protease which then cleaves and activates effector caspases caspase-3 (CASP3) or caspase-7 (CASP7). Promotes DNA damage- induced apoptosis in a ABL1/c-Abl-dependent manner. Proteolytically cleaves poly(ADP-ribose) polymerase (PARP). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:<a href="http://www.uniprot.org/citations/36758105" target="\_blank">36758105</a>, PubMed:<a href="http://www.uniprot.org/citations/36758106" target="\_blank">36758106</a>).

### Tissue Location


Ubiquitous, with highest expression in the heart, moderate expression in liver, skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart, specifically expressed in myocytes.


## Anti-Caspase-9 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Caspase-9 Picoband Antibody - Images





Anti- Caspase-9 Picoband antibody, ABO12023, Western blotting All lanes: Anti Caspase-9 (ABO12023) at 0.5ug/ml Lane 1: A549 Whole Cell Lysate at 40ug Lane 2: SMMC Whole Cell Lysate at 40ug Lane 3: 293T Whole Cell Lysate at 40ug Lane 4: JURKAT Whole Cell Lysate at 40ug Lane 5: RAJI Whole Cell Lysate at 40ug Lane 6: CEM Whole Cell Lysate at 40ug Lane 7: HUT Whole Cell Lysate at 40ug Predicted bind size: 35KD Observed bind size: 35KD

#### **Anti-Caspase-9 Picoband Antibody - Background**

CASP9 is also known as MCH6 or APAF3. This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein can undergo autoproteolytic processing and activation by the apoptosome, a protein complex of cytochrome c and the apoptotic peptidase activating factor 1; this step is thought to be one of the earliest in the caspase activation cascade. This protein is thought to play a central role in apoptosis and to be a tumor suppressor. Alternative splicing results in multiple transcript variants.