

Anti-SFRP2 Picoband Antibody
Catalog # ABO12053**Specification****Anti-SFRP2 Picoband Antibody - Product Information**

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q96HF1 |
| Host | Rabbit |
| Reactivity | Human |
| Clonality | Polyclonal |
| Format | Lyophilized |

Description

Rabbit IgG polyclonal antibody for Secreted frizzled-related protein 2(SFRP2) detection. Tested with WB, ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SFRP2 Picoband Antibody - Additional Information

Gene ID 6423

Other Names

Secreted frizzled-related protein 2, FRP-2, sFRP-2, Secreted apoptosis-related protein 1, SARP-1, SFRP2, FRP2, SARP1

Calculated MW

33490 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, -
ELISA , 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted .

Tissue Specificity

Expressed in adipose tissue, heart, brain, skeletal muscle, pancreas, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in adipose tissue, small intestine and colon. .

Protein Name

Secreted frizzled-related protein 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E.coli-derived human SFRP2 recombinant protein (Position: L104-C295). Human SFRP2 shares 99% amino acid (aa) sequence identity with mouse SFRP2.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the secreted frizzled-related protein (sFRP) family.

Anti-SFRP2 Picoband Antibody - Protein Information

Name SFRP2

Synonyms FRP2, SARP1

Function

Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types. SFRP2 may be important for eye retinal development and for myogenesis.

Cellular Location

Secreted.

Tissue Location


Expressed in adipose tissue, heart, brain, skeletal muscle, pancreas, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in adipose tissue, small intestine and colon


Anti-SFRP2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SFRP2 Picoband Antibody - Images





Anti- SFRP2 Picoband antibody, ABO12053, Western blottingAll lanes: Anti SFRP2 (ABO12053) at 0.5ug/mlLane 1: COLO320 Whole Cell Lysate at 40ugLane 2: SW620 Whole Cell Lysate at 40ugPredicted bind size: 33KDObserved bind size: 33KD

Anti-SFRP2 Picoband Antibody - Background

SFRP2 is also known as FRP-2, SARP1 or SDF-5. It is Expressed in adipose tissue, heart, brain, skeletal muscle, pancreas, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in adipose tissue, small intestine and colon. This gene encodes a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. Methylation of this gene is a potential marker for the presence of colorectal cancer.Â