

Anti-ADRB2 Picoband Antibody
Catalog # ABO12056**Specification****Anti-ADRB2 Picoband Antibody - Product Information**

Application	WB
Primary Accession	P07550
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Beta-2 adrenergic receptor(ADRB2) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-ADRB2 Picoband Antibody - Additional Information

Gene ID 154

Other Names

Beta-2 adrenergic receptor, Beta-2 adrenoreceptor, Beta-2 adrenoceptor, ADRB2, ADRB2R, B2AR

Calculated MW

46459 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Rat, Human

Subcellular Localization

Cell membrane ; Multi-pass membrane protein . Colocalizes with VHL at the cell membrane.

Protein Name

Beta-2 adrenergic receptor

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human ADRB2(221-256aa RVFQEAKRQLQKIDKSEGRFHVQNLSQVEQDGRGTGH), different from the related mouse and rat sequences by three amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the G-protein coupled receptor 1 family. Adrenergic receptor subfamily. ADRB2 sub-subfamily.

Anti-ADRB2 Picoband Antibody - Protein Information

Name ADRB2

Synonyms ADRB2R, B2AR

Function

Beta-adrenergic receptors mediate the catecholamine-induced activation of adenylate cyclase through the action of G proteins. The beta-2-adrenergic receptor binds epinephrine with an approximately 30- fold greater affinity than it does norepinephrine.

Cellular Location

Cell membrane; Multi-pass membrane protein. Early endosome. Golgi apparatus. Note=Colocalizes with VHL at the cell membrane (PubMed:19584355). Activated receptors are internalized into endosomes prior to their degradation in lysosomes (PubMed:20559325) Activated receptors are also detected within the Golgi apparatus (PubMed:27481942).


Anti-ADRB2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ADRB2 Picoband Antibody - Images





Anti- ADRB2 Picoband antibody, ABO12056, Western blotting All lanes: AntiADRB2 (ABO12056) at 0.5ug/ml WB: Rat Brain Tissue Lysate at 50ug Predicted bind size: 47KD Observed bind size: 47KD

Anti-ADRB2 Picoband Antibody - Background

ADRB2, also known as beta-2 adrenergic receptor, is a beta-adrenergic receptor within a cell membrane which reacts with adrenaline (epinephrine) as a hormone or neurotransmitter affecting muscles or organs. It is mapped to 5q32. This receptor is directly associated with one of its ultimate effectors, the class C L-type calcium channel Ca(V)1.2. The genetic variation in the ADRB2 gene may be of major importance for obesity, energy expenditure, and lipolytic ADRB2 function in adipose tissue, at least in women. What's more, it has been found that activation of ADRB2 receptors can stimulate gamma-secretase activity and beta-amyloid production, and the ADRB2 receptors activator may contribute to beta-amyloid accumulation in AD.