

**Anti-Cyclin A1 Picoband Antibody**  
Catalog # ABO12173**Specification**

---

**Anti-Cyclin A1 Picoband Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P78396</a>
Host	<b>Rabbit</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Format	<b>Lyophilized</b>

**Description**

Rabbit IgG polyclonal antibody for Cyclin-A1(CCNA1) detection. Tested with WB in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Cyclin A1 Picoband Antibody - Additional Information**

**Gene ID** 8900

**Other Names**

Cyclin-A1, CCNA1

**Calculated MW**

52358 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Nucleus .

**Tissue Specificity**

Very high levels in testis and very low levels in brain. Also found in myeloid leukemia cell lines.

**Protein Name**

Cyclin-A1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>N.

**Immunogen**

E.coli-derived human Cyclin A1 recombinant protein (Position: D64-E263). Human Cyclin A1 shares 78.5% and 79.5% amino acid (aa) sequence identity with mouse and rat Cyclin A1, respectively.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

### Sequence Similarities

Belongs to the cyclin family. Cyclin AB subfamily.

## Anti-Cyclin A1 Picoband Antibody - Protein Information

**Name** CCNA1

### Function

May be involved in the control of the cell cycle at the G1/S (start) and G2/M (mitosis) transitions. May primarily function in the control of the germline meiotic cell cycle and additionally in the control of mitotic cell cycle in some somatic cells.

### Cellular Location

Nucleus.

### Tissue Location


Very high levels in testis and very low levels in brain. Also found in myeloid leukemia cell lines


## Anti-Cyclin A1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Cyclin A1 Picoband Antibody - Images





Anti- Cyclin A1 Picoband antibody, ABO12173, Western blotting All lanes: Anti Cyclin A1 (ABO12173) at 0.5ug/ml WB: Human Placenta Tissue Lysate at 50ug Predicted bind size: 52KD Observed bind size: 52KD

#### **Anti-Cyclin A1 Picoband Antibody - Background**

Cyclin-A1 is a protein that in humans is encoded by the CCNA1 gene. The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. The cyclin encoded by this gene was shown to be expressed in testis and brain, as well as in several leukemic cell lines, and is thought to primarily function in the control of the germline meiotic cell cycle. This cyclin binds both CDK2 and CDC2 kinases, which give two distinct kinase activities, one appearing in S phase, the other in G2, and thus regulate separate functions in cell cycle. Also this cyclin was found to bind to important cell cycle regulators, such as Rb family proteins, transcription factor E2F-1, and the p21 family proteins. Multiple transcript variants encoding different isoforms have been found for this gene.