

**Anti-NQO1 Picoband Antibody**  
Catalog # ABO12185**Specification****Anti-NQO1 Picoband Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P15559</a>
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for NAD(P)H dehydrogenase [quinone] 1(NQO1) detection. Tested with WB in Human;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-NQO1 Picoband Antibody - Additional Information**

**Gene ID** 1728

**Other Names**

NAD(P)H dehydrogenase [quinone] 1, 1.6.5.2, Azoreductase, DT-diaphorase, DTD, Menadione reductase, NAD(P)H:quinone oxidoreductase 1, Phylloquinone reductase, Quinone reductase 1, QR1, NQO1, DIA4, NMOR1

**Calculated MW**

30868 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Rat<br>

**Subcellular Localization**

Cytoplasm.

**Protein Name**

NAD(P)H dehydrogenase [quinone] 1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>N.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human NQO1 (242-274aa EVQDEEKNNKFGLSVGHHLGKSIPTDNQIKARK), different from the related mouse and rat sequences by five amino acids.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

### Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

### Sequence Similarities

Belongs to the NAD(P)H dehydrogenase (quinone) family.

## Anti-NQO1 Picoband Antibody - Protein Information

**Name** NQO1 {ECO:0000303|PubMed:1657151, ECO:0000312|HGNC:HGNC:2874}

### Function

Flavin-containing quinone reductase that catalyzes two- electron reduction of quinones to hydroquinones using either NADH or NADPH as electron donors. In a ping-pong kinetic mechanism, the electrons are sequentially transferred from NAD(P)H to flavin cofactor and then from reduced flavin to the quinone, bypassing the formation of semiquinone and reactive oxygen species (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/8999809" target="\_blank">8999809</a>, PubMed:<a href="http://www.uniprot.org/citations/9271353" target="\_blank">9271353</a>). Regulates cellular redox state primarily through quinone detoxification. Reduces components of plasma membrane redox system such as coenzyme Q and vitamin quinones, producing antioxidant hydroquinone forms. In the process may function as superoxide scavenger to prevent hydroquinone oxidation and facilitate excretion (PubMed:<a href="http://www.uniprot.org/citations/15102952" target="\_blank">15102952</a>, PubMed:<a href="http://www.uniprot.org/citations/8999809" target="\_blank">8999809</a>, PubMed:<a href="http://www.uniprot.org/citations/9271353" target="\_blank">9271353</a>). Alternatively, can activate quinones and their derivatives by generating redox reactive hydroquinones with DNA cross-linking antitumor potential (PubMed:<a href="http://www.uniprot.org/citations/8999809" target="\_blank">8999809</a>). Acts as a gatekeeper of the core 20S proteasome known to degrade proteins with unstructured regions. Upon oxidative stress, interacts with tumor suppressors TP53 and TP73 in a NADH-dependent way and inhibits their ubiquitin-independent degradation by the 20S proteasome (PubMed:<a href="http://www.uniprot.org/citations/15687255" target="\_blank">15687255</a>, PubMed:<a href="http://www.uniprot.org/citations/28291250" target="\_blank">28291250</a>).

### Cellular Location


Cytoplasm, cytosol {ECO:0000250|UniProtKB:P05982}

## Anti-NQO1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-NQO1 Picoband Antibody - Images



Anti- NQO1 Picoband antibody, ABO12185, Western blottingAll lanes: Anti NQO1 (ABO12185) at 0.5ug/mlLane 1: Rat Liver Tissue Lysate at 50ugLane 2: Rat Lung Tissue Lysate at 50ugLane 3: HELA Whole Cell Lysate at 40ugLane 4: A549 Whole Cell Lysate at 40ugLane 5: MM231 Whole Cell Lysate at 40ugLane 6: SW620 Whole Cell Lysate at 40ugLane 7: 22RV1 Whole Cell Lysate at 40ugPredicted bind size: 31KDObserved bind size: 31KD

## Anti-NQO1 Picoband Antibody - Background

This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. And this FAD-binding protein forms homodimers and reduces quinones to hydroquinones. In addition, this protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized.