

**Anti-Stathmin 1 Picoband Antibody**  
Catalog # ABO12247**Specification****Anti-Stathmin 1 Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P16949</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Stathmin(STMN1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Stathmin 1 Picoband Antibody - Additional Information**

Gene ID 3925

**Other Names**

Stathmin, Leukemia-associated phosphoprotein p18, Metablastin, Oncoprotein 18, Op18, Phosphoprotein p19, pp19, Prosolin, Protein Pr22, pp17, STMN1, C1orf215, LAP18, OP18

**Calculated MW**

17303 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat<br>Western blot, 0.1-0.5 µg/ml, Human, Mouse<br>

**Subcellular Localization**

Cytoplasm, cytoskeleton.

**Tissue Specificity**

Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients with chronic lymphoid or myeloid leukemia. .

**Protein Name**

Stathmin

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

#### Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human Stathmin 1 (2-34aa ASSDIQVKELEKRASGQAFELILSPRSKESVPE), different from the related mouse sequence by one amino acid, and identical to the related rat sequence.

#### Purification

Immunogen affinity purified.

#### Cross Reactivity

No cross reactivity with other proteins

#### Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

#### Sequence Similarities

Belongs to the stathmin family.

### Anti-Stathmin 1 Picoband Antibody - Protein Information

**Name** STMN1

**Synonyms** C1orf215, LAP18, OP18

#### Function

Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser-16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).

#### Cellular Location

Cytoplasm, cytoskeleton.

#### Tissue Location

Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients with chronic lymphoid or myeloid leukemia.


### Anti-Stathmin 1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-Stathmin 1 Picoband Antibody - Images




Anti- Stathmin 1 Picoband antibody, ABO12247, Western blottingAll lanes: Anti Stathmin 1 (ABO12247) at 0.5ug/mlLane 1: Mouse Brain Tissue Lysate at 50ugLane 2: Mouse Testis Tissue Lysate at 50ugLane 3: MM231 Whole Cell Lysate at 40ugLane 4: MCF-7 Whole Cell Lysate at 40ugPredicted bind size: 17KDObserved bind size: 17KD



Anti- Stathmin 1 Picoband antibody, ABO12247, IHC(P)IHC(P): Mouse Testis Tissue



anti- Stathmin 1 Picoband antibody, ABO12247, IHC(P)IHC(P): Rat Brain Tissue



anti- Stathmin 1 Picoband antibody, ABO12247, IHC(P)IHC(P): Human Mammary Cancer Tissue

### **Anti-Stathmin 1 Picoband Antibody - Background**

Stathmin 1/oncoprotein 18, also known as STMN1, is a highly conserved 17 kDa protein. This gene belongs to the stathmin family of genes. It encodes a ubiquitous cytosolic phosphoprotein proposed to function as an intracellular relay integrating regulatory signals of the cellular environment. The encoded protein is involved in the regulation of the microtubule filament system by destabilizing microtubules. It prevents assembly and promotes disassembly of microtubules. Multiple transcript variants encoding different isoforms have been found for this gene.