

Anti-MAOA Picoband Antibody
Catalog # ABO12350

Specification

Anti-MAOA Picoband Antibody - Product Information

Application	WB, IHC
Primary Accession	P21397
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Amine oxidase [flavin-containing] A(MAOA) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MAOA Picoband Antibody - Additional Information

Gene ID 4128

Other Names

Amine oxidase [flavin-containing] A, 1.4.3.4, Monoamine oxidase type A, MAO-A, MAOA

Calculated MW

59682 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Mitochondrion outer membrane; Single-pass type IV membrane protein; Cytoplasmic side.

Tissue Specificity

Heart, liver, duodenum, blood vessels and kidney.

Protein Name

Amine oxidase [flavin-containing] A

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human MAOA (457-493aa REVLNGLGKVTEDIWVQEPESKDVPAVEITHTFWER), different from the related mouse sequence by five amino acids, and from the related rat sequence by six amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-MAOA Picoband Antibody - Protein Information

Name MAOA ([HGNC:6833](#))

Function

Catalyzes the oxidative deamination of primary and some secondary amine such as neurotransmitters, with concomitant reduction of oxygen to hydrogen peroxide and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed: [18391214](http://www.uniprot.org/citations/18391214) target="_blank">18391214, PubMed: [20493079](http://www.uniprot.org/citations/20493079) target="_blank">20493079, PubMed: [24169519](http://www.uniprot.org/citations/24169519) target="_blank">24169519, PubMed: [8316221](http://www.uniprot.org/citations/8316221) target="_blank">8316221). Preferentially oxidizes serotonin (PubMed: [20493079](http://www.uniprot.org/citations/20493079) target="_blank">20493079, PubMed: [24169519](http://www.uniprot.org/citations/24169519) target="_blank">24169519). Also catalyzes the oxidative deamination of kynuramine to 3-(2-aminophenyl)-3-oxopropanal that can spontaneously condense to 4-hydroxyquinoline (By similarity).

Cellular Location

Mitochondrion outer membrane {ECO:0000250|UniProtKB:P21396}; Single-pass type IV membrane protein {ECO:0000250|UniProtKB:P21396}; Cytoplasmic side {ECO:0000250|UniProtKB:P21396}

Tissue Location


Heart, liver, duodenum, blood vessels and kidney.


Anti-MAOA Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MAOA Picoband Antibody - Images







Anti- MAOA Picoband antibody, ABO12350, Western blotting All lanes: Anti MAOA (ABO12350) at 0.5ug/ml Lane 1: Rat Kidney Tissue Lysate at 50ug Lane 2: Mouse Kidney Tissue Lysate at 50ug Lane 3: COLO320 Whole Cell Lysate at 40ug Lane 4: HEPG2 Whole Cell Lysate at 40ug Lane 5: HEPA Whole Cell Lysate at 40ug Predicted bind size: 60KD Observed bind size: 60KD



Anti- MAOA Picoband antibody, ABO12350, IHC(P) IHC(P): Mouse Cardiac Muscle Tissue



Anti- MAOA Picoband antibody, ABO12350, IHC(P) IHC(P): Rat Cardiac Muscle Tissue



Anti- MAOA Picoband antibody, ABO12350,IHC(P)IHC(P): Human Intestinal Cancer Tissue

Anti-MAOA Picoband Antibody - Background

MAOA(Monoamine oxidase A), also known as AMINE OXIDASE (FLAVIN-CONTAINING) A, is an enzyme that in humans is encoded by the MAO-A gene. MAOA is an isozyme of monoamine oxidase which is also mapped on Xp11.3. MAOA degrades amine neurotransmitters, such as dopamine, norepinephrine, and serotonin. The protein localizes to the outer mitochondrial membrane. Mutation in MAOA results in monoamine oxidase deficiency, or Brunner syndrome. In humans, there is a 30-base repeat sequence repeated in one of several different numbers of times in the promoter region of the gene coding for MAOA. MAO-A levels in the brain as measured using positron emission tomography are elevated by an average of 34% in patients with major depressive disorder. Inhibition of MAOA prevented apoptosis, and serum starvation of cortical brain cells from Maa-deficient mice resulted in reduced apoptosis compared with wildtype mice.