

**Anti-NDRG2 Picoband Antibody**  
Catalog # ABO12358

**Specification**

---

**Anti-NDRG2 Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q9UN36</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Protein NDRG2(NDRG2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-NDRG2 Picoband Antibody - Additional Information**

**Gene ID** 57447

**Other Names**

Protein NDRG2, N-myc downstream-regulated gene 2 protein, Protein Syld709613, NDRG2, KIAA1248, SYLD

**Calculated MW**

40798 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, Rat, Human, By Heat  
Western blot, 0.1-0.5 µg/ml, Mouse, Rat, Human

**Subcellular Localization**

Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone . In neurons, seems to concentrate at axonal growth cone. Perinuclear in neurons (By similarity) .

**Tissue Specificity**

Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Expression levels are low in pancreatic and liver cancer tissues; absent in meningioma. Expressed in low-grade gliomas but present at low levels in glioblastoma. Isoform 1 and isoform 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein level) .

**Protein Name**

Protein NDRG2

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human NDRG2 (210-247aa NSELIQKYRNIITHAPNLDNIELYWNSYNNRRDLNFER), different from the related mouse and rat sequences by three amino acids.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Anti-NDRG2 Picoband Antibody - Protein Information**

**Name** NDRG2

**Synonyms** KIAA1248, SYLD

**Function**

Contributes to the regulation of the Wnt signaling pathway. Down-regulates CTNNB1-mediated transcriptional activation of target genes, such as CCND1, and may thereby act as tumor suppressor. May be involved in dendritic cell and neuron differentiation.

**Cellular Location**

Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone. Note=In neurons, seems to concentrate at axonal growth cone. Perinuclear in neurons (By similarity).


**Tissue Location**


Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Expression levels are low in pancreatic and liver cancer tissues; absent in meningioma Expressed in low-grade gliomas but present at low levels in glioblastoma. Isoform 1 and isoform 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein level)

**Anti-NDRG2 Picoband Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Anti-NDRG2 Picoband Antibody - Images**



Anti- NDRG2 Picoband antibody, ABO12358, Western blotting All lanes: Anti NDRG2 (ABO12358) at 0.5ug/ml Lane 1: Rat Brain Tissue Lysate at 50ug Lane 2: Mouse Brain Tissue Lysate at 50ug Predicted bind size: 41KD Observed bind size: 41KD



Anti- NDRG2 Picoband antibody, ABO12358, IHC(P) IHC(P): Mouse Brain Tissue



Anti- NDRG2 Picoband antibody, ABO12358, IHC(P) IHC(P): Rat Brain Tissue

**Anti-NDRG2 Picoband Antibody - Background**

Protein NDRG2, also known as KIAA1248, is a protein that in humans is encoded by the NDRG2 gene. It is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. This gene is mapped to 14q11.2. The protein encoded by this gene is a cytoplasmic protein that may play a role in neurite outgrowth. This gene may be involved in

glioblastoma carcinogenesis. It contributes to the regulation of the WNT signaling pathway. Also, this gene may be involved in dendritic cell and neuron differentiation.