

Anti-SHP2 Picoband Antibody
Catalog # ABO12361**Specification****Anti-SHP2 Picoband Antibody - Product Information**

| | |
|-------------------|------------------------|
| Application | WB, IHC |
| Primary Accession | Q06124 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Clonality | Polyclonal |
| Format | Lyophilized |

Description

Rabbit IgG polyclonal antibody for Tyrosine-protein phosphatase non-receptor type 11(PTPN11) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SHP2 Picoband Antibody - Additional Information

Gene ID 5781

Other Names

Tyrosine-protein phosphatase non-receptor type 11, 3.1.3.48, Protein-tyrosine phosphatase 1D, PTP-1D, Protein-tyrosine phosphatase 2C, PTP-2C, SH-PTP2, SHP-2, Shp2, SH-PTP3, PTPN11, PTP2C, SHPTP2

Calculated MW

68436 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cytoplasm.

Tissue Specificity

Widely expressed, with highest levels in heart, brain, and skeletal muscle. .

Protein Name

Tyrosine-protein phosphatase non-receptor type 11

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human SHP2 (69-99aa EK_FATLAELVQYYMEHHGQLKEKNGDVIELK), identical to the related mouse and rat sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-SHP2 Picoband Antibody - Protein Information

Name PTPN11

Synonyms PTP2C, SHPTP2

Function

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:10655584, PubMed:18559669, PubMed:18829466, PubMed:26742426, PubMed:28074573). Positively regulates MAPK signal transduction pathway (PubMed:28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:18559669). Dephosphorylates CDC73 (PubMed:26742426). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:19275884).

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Widely expressed, with highest levels in heart, brain, and skeletal muscle.


Anti-SHP2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-SHP2 Picoband Antibody - Images




Anti- SHP2 Picoband antibody, ABO12361, Western blotting All lanes: Anti SHP2 (ABO12361) at 0.5ug/ml Lane 1: Rat Brain Tissue Lysate at 50ug Lane 2: Mouse Brain Tissue Lysate at 50ug Lane 3: Rat Cardiac Muscle Tissue Lysate at 50ug Lane 4: Mouse Kidney Tissue Lysate at 50ug Lane 5: HELA Whole Cell Lysate at 40ug Lane 6: SW620 Whole Cell Lysate at 40ug Lane 7: HEPG2 Whole Cell Lysate at 40ug Lane 8: JURKAT Whole Cell Lysate at 40ug Predicted bind size: 68KD Observed bind size: 68KD



Anti- SHP2 Picoband antibody, ABO12361, IHC(P) IHC(P): Mouse Cardiac Muscle Tissue



Anti- SHP2 Picoband antibody, ABO12361,IHC(P)IHC(P): Rat Skeletal Muscle Tissue



Anti- SHP2 Picoband antibody, ABO12361,IHC(P)IHC(P): Human Lung Cancer Tissue

Anti-SHP2 Picoband Antibody - Background

PTPN11 (Tyrosine-protein phosphatase non-receptor type 11), also known as protein-tyrosine phosphatase 1D (PTP-1D), protein-tyrosine phosphatase 2C (PTP-2C), TYROSINE PHOSPHATASE SHP2 (SHP2), BPTP3, SH-PTP2, SHP-2, SH-PTP3, is an enzyme that in humans is encoded by the PTPN11 gene. PTPN11 is a member of the protein tyrosine phosphatase (PTP) family. The open reading frame consists of 1,779 nucleotides potentially encoding a protein of 593 amino acids with a predicted molecular mass of 68 kD. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates. This PTP is widely expressed in most tissues and plays a regulatory role in various cell signaling events that are important for a diversity of cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration. Mutations in this gene are a cause of Noonan syndrome as well as acute myeloid leukemia.