

Anti-LOXL2 Picoband Antibody
Catalog # ABO12406**Specification****Anti-LOXL2 Picoband Antibody - Product Information**

Application	WB
Primary Accession	Q9Y4K0
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Lysyl oxidase homolog 2(LOXL2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-LOXL2 Picoband Antibody - Additional Information

Gene ID 4017

Other Names

Lysyl oxidase homolog 2, 1.4.3.13, Lysyl oxidase-like protein 2, Lysyl oxidase-related protein 2, Lysyl oxidase-related protein WS9-14, LOXL2

Calculated MW

86725 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Secreted, extracellular space, extracellular matrix, basement membrane . Nucleus. Chromosome. Associated with chromatin. It is unclear how LOXL2 is nuclear: it contains a clear signal sequence and is predicted to localize in the extracellular medium. However, different reports confirmed the intracellular location and its key role in transcription regulation.

Tissue Specificity

Expressed in many tissues. Highest expression in reproductive tissues, placenta, uterus and prostate. Up- regulated in a number of cancers cells and tissues.

Protein Name

Lysyl oxidase homolog 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human LOXL2 (739-770aa HRIWMYNCHIGGSFSEETEKKFEHFSGLLNQ), different from the related mouse and rat sequences by four amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-LOXL2 Picoband Antibody - Protein Information

Name LOXL2

Function

Mediates the post-translational oxidative deamination of lysine residues on target proteins leading to the formation of deaminated lysine (allysine) (PubMed: [27735137](http://www.uniprot.org/citations/27735137)). Acts as a transcription corepressor and specifically mediates deamination of trimethylated 'Lys-4' of histone H3 (H3K4me3), a specific tag for epigenetic transcriptional activation (PubMed: [27735137](http://www.uniprot.org/citations/27735137)). Shows no activity against histone H3 when it is trimethylated on 'Lys-9' (H3K9me3) or 'Lys-27' (H3K27me3) or when 'Lys-4' is monomethylated (H3K4me1) or dimethylated (H3K4me2) (PubMed: [27735137](http://www.uniprot.org/citations/27735137)). Also mediates deamination of methylated TAF10, a member of the transcription factor IID (TFIID) complex, which induces release of TAF10 from promoters, leading to inhibition of TFIID-dependent transcription (PubMed: [25959397](http://www.uniprot.org/citations/25959397)). LOXL2-mediated deamination of TAF10 results in transcriptional repression of genes required for embryonic stem cell pluripotency including POU5F1/OCT4, NANOG, KLF4 and SOX2 (By similarity). Involved in epithelial to mesenchymal transition (EMT) via interaction with SNAI1 and participates in repression of E-cadherin CDH1, probably by mediating deamination of histone H3 (PubMed: [16096638](http://www.uniprot.org/citations/16096638), PubMed: [24414204](http://www.uniprot.org/citations/24414204), PubMed: [27735137](http://www.uniprot.org/citations/27735137)). During EMT, involved with SNAI1 in negatively regulating pericentromeric heterochromatin transcription (PubMed: [24239292](http://www.uniprot.org/citations/24239292)). SNAI1 recruits LOXL2 to pericentromeric regions to oxidize histone H3 and repress transcription which leads to release of heterochromatin component CBX5/HP1A, enabling chromatin reorganization and acquisition of mesenchymal traits (PubMed: [24239292](http://www.uniprot.org/citations/24239292)). Interacts with the endoplasmic reticulum protein HSPA5 which activates the IRE1-XBP1 pathway of the unfolded protein response, leading to expression of several transcription factors involved in EMT and subsequent EMT induction (PubMed: [28332555](http://www.uniprot.org/citations/28332555)). Involved in E-cadherin repression following hypoxia, a hallmark of EMT believed to amplify tumor aggressiveness, suggesting that it may play a role in tumor progression (PubMed: [20026874](http://www.uniprot.org/citations/20026874)). When secreted into the extracellular matrix, promotes cross-linking of extracellular matrix proteins by mediating oxidative deamination of peptidyl lysine residues in precursors to fibrous collagen and elastin (PubMed: [20026874](http://www.uniprot.org/citations/20026874)).

[20306300](http://www.uniprot.org/citations/20306300)). Acts as a regulator of sprouting angiogenesis, probably via collagen IV scaffolding (PubMed:[21835952](http://www.uniprot.org/citations/21835952)). Acts as a regulator of chondrocyte differentiation, probably by regulating expression of factors that control chondrocyte differentiation (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix, basement membrane. Nucleus. Chromosome. Endoplasmic reticulum. Note=Associated with chromatin (PubMed:27735137). It is unclear how LOXL2 is nuclear as it contains a signal sequence and has been shown to be secreted (PubMed:23319596) However, a number of reports confirm its intracellular location and its key role in transcription regulation (PubMed:22204712, PubMed:22483618).

Tissue Location


Expressed in many tissues (PubMed:10212285). Highest expression in reproductive tissues, placenta, uterus and prostate (PubMed:10212285). In esophageal epithelium, expressed in the basal, prickle and granular cell layers (PubMed:22204712). Up-regulated in a number of cancers cells and tissues.

Anti-LOXL2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-LOXL2 Picoband Antibody - Images



Anti- LOXL2 Picoband antibody, ABO12406, Western blottingAll lanes: Anti LOXL2 (ABO12406) at 0.5ug/mlLane 1: Mouse Testis Tissue Lysate at 50ugLane 2: Rat Ovary Tissue Lysate at 50ugLane 3: Human Placenta Tissue Lysate at 50ugLane 4: HELA Whole Cell Lysate at 40ugLane 5: 22RV1

Whole Cell Lysate at 40ugLane 6: MCF-7 Whole Cell Lysate at 40ugPredicted bind size:
87KDObserved bind size: 87KD

Anti-LOXL2 Picoband Antibody - Background

Lysyl oxidase homolog 2 is an enzyme that in humans is encoded by the LOXL2 gene. This gene encodes a member of the lysyl oxidase gene family. The prototypic member of the family is essential to the biogenesis of connective tissue, encoding an extracellular copper-dependent amine oxidase that catalyses the first step in the formation of crosslinks in collagens and elastin. A highly conserved amino acid sequence at the C-terminus end appears to be sufficient for amine oxidase activity, suggesting that each family member may retain this function. The N-terminus is poorly conserved and may impart additional roles in developmental regulation, senescence, tumor suppression, cell growth control, and chemotaxis to each member of the family. LOXL2 can also crosslink collagen type IV and hence influence the sprouting of new blood vessels.