

Anti-ROC1 Picoband Antibody

Catalog # ABO12484

Specification

Anti-ROC1 Picoband Antibody - Product Information

ApplicationWBPrimary AccessionP62877HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for E3 ubiquitin-protein ligase RBX1(RBX1) detection. Tested withWB in Human;Mouse;Rat.WB

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-ROC1 Picoband Antibody - Additional Information

Gene ID 9978

Other Names E3 ubiquitin-protein ligase RBX1, 6.3.2.-, Protein ZYP, RING finger protein 75, RING-box protein 1, Rbx1, Regulator of cullins 1, E3 ubiquitin-protein ligase RBX1, N-terminally processed, RBX1, RNF75, ROC1

Calculated MW 12274 MW KDa

Application Details Western blot, 0.1-0.5 μg/ml, Mouse, Rat, Human

Subcellular Localization Cytoplasm . Nucleus .

Tissue Specificity Widely expressed.

Protein Name E3 ubiquitin-protein ligase RBX1

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human ROC1 (76-108aa NHAFHFHCISRWLKTRQVCPLDNREWEFQKYGH), identical to the related mouse sequence.



Purification Immunogen affinity purified.

Cross Reactivity No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-ROC1 Picoband Antibody - Protein Information

Name RBX1 (HGNC:9928)

Function

E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase (CRLs) complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair (PubMed:10230407, PubMed:10579999, PubMed:11961546, PubMed:15983046, PubMed:16678110, PubMed:19112177, PubMed:19679664, PubMed:22748924, PubMed:23455478, PubMed:27565346, PubMed:29769719, PubMed:33417871, PubMed:38326650). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiguitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (PubMed:27565346). The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components. As a component of the CSA complex promotes the ubiguitination of ERCC6 resulting in proteasomal degradation. Core component of the Cul7-RING(FBXW8) ubiquitin ligase complex, which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:35982156). Core component of a Cul9-RING ubiquitin ligase complex composed of CUL9 and RBX1, which mediates mono-ubiquitination of p53/TP53 (PubMed: 38605244). Recruits the E2 ubiguitin-conjugating enzyme CDC34 to the complex and brings it into close proximity to the substrate. Probably also stimulates CDC34 autoubiquitination. May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair. Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M. Involved in the ubiquitination of KEAP1, ENC1 and KLHL41. In concert with ATF2 and CUL3, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. As part of a multisubunit complex composed of elongin BC complex (ELOB and ELOC), elongin A/ELOA, RBX1 and CUL5; polyubiquitinates monoubiquitinated POLR2A (PubMed:19920177).

Cellular Location



Cytoplasm. Nucleus

Tissue Location Widely expressed.

Anti-ROC1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-ROC1 Picoband Antibody - Images



Anti- ROC1 Picoband antibody, ABO12484, Western blottingAll lanes: Anti ROC1 (ABO12484) at 0.5ug/mlLane 1: Rat Testis Tissue Lysate at 50ugLane 2: Rat Brain Tissue Lysate at 50ugLane 3: Mouse Brain Tissue Lysate at 50ugLane 4: Mouse Spleen Tissue Lysate at 50ugPredicted bind size: 15KDObserved bind size: 15KD

Anti-ROC1 Picoband Antibody - Background

RING-box protein 1, also known as ROC1, is a protein that in humans is encoded by the RBX1 gene. This gene is mapped to chromosome 22q13.2 based on an alignment of the RBX1 sequence with the genomic sequence. ROC1 is recruited by cullin-1 to form a quaternary SCF (HOS)-ROC1 holoenzyme (with SKP1 and the BTRCP homolog HOS). SCF (HOS)-ROC1 binds IKK-beta-phosphorylated I-kappa-B-alpha and catalyzes its ubiquitination in the presence of ubiquitin, E1, and CDC34. Conclusively, ROC1 plays a unique role in the ubiquitination reaction by heterodimerizing with cullin-1 to catalyze ubiquitin polymerization.