

Anti-TIF1 Gamma Picoband Antibody
Catalog # ABO12522**Specification****Anti-TIF1 Gamma Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	Q9UPN9
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for E3 ubiquitin-protein ligase TRIM33(TRIM33) detection. Tested with WB, IHC-P in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-TIF1 Gamma Picoband Antibody - Additional Information

Gene ID 51592

Other Names

E3 ubiquitin-protein ligase TRIM33, 6.3.2.-, Ectodermin homolog, RET-fused gene 7 protein, Protein Rfg7, Transcription intermediary factor 1-gamma, TIF1-gamma, Tripartite motif-containing protein 33, TRIM33, KIAA1113, RFG7, TIF1G

Calculated MW

122533 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Nucleus . In discrete nuclear dots resembling nuclear bodies. .

Tissue Specificity

Expressed in stem cells at the bottom of the crypts of the colon (at protein level). Expressed in colon adenomas and adenocarcinomas (at protein level). Expressed in brain, lung, liver, spleen, thymus, prostate, kidney, testis, heart, placenta, pancreas, small intestine, ovary, colon, skeletal muscle and hematopoietic progenitors.

Protein Name

E3 ubiquitin-protein ligase TRIM33

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E.coli-derived human TIF1 gamma recombinant protein (Position: M1001-K1127). Human TIF1 gamma shares 96.1% amino acid (aa) sequence identity with mouse TIF1 gamma.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-TIF1 Gamma Picoband Antibody - Protein Information

Name TRIM33

Synonyms KIAA1113, RFG7, TIF1G

Function

Acts as an E3 ubiquitin-protein ligase. Promotes SMAD4 ubiquitination, nuclear exclusion and degradation via the ubiquitin proteasome pathway. According to PubMed: [16751102](http://www.uniprot.org/citations/16751102), does not promote a decrease in the level of endogenous SMAD4. May act as a transcriptional repressor. Inhibits the transcriptional response to TGF-beta/BMP signaling cascade. Plays a role in the control of cell proliferation. Its association with SMAD2 and SMAD3 stimulates erythroid differentiation of hematopoietic stem/progenitor (By similarity). Monoubiquitinates SMAD4 and acts as an inhibitor of SMAD4-dependent TGF-beta/BMP signaling cascade (Monoubiquitination of SMAD4 hampers its ability to form a stable complex with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade).

Cellular Location

Nucleus. Note=In discrete nuclear dots resembling nuclear bodies (By similarity). Localizes to sites of DNA damage (PubMed:25593309). {ECO:0000250|UniProtKB:Q99PP7, ECO:0000269|PubMed:25593309}

Tissue Location

Expressed in stem cells at the bottom of the crypts of the colon (at protein level). Expressed in colon adenomas and adenocarcinomas (at protein level). Expressed in brain, lung, liver, spleen, thymus, prostate, kidney, testis, heart, placenta, pancreas, small intestine, ovary, colon, skeletal muscle and hematopoietic progenitors


Anti-TIF1 Gamma Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TIF1 Gamma Picoband Antibody - Images



Anti- TIF1 gamma Picoband antibody, ABO12522, Western blotting
All lanes: Anti TIF1 gamma (ABO12522) at 0.5ug/ml
Lane 1: 22RV1 Whole Cell Lysate at 40ug
Lane 2: SW620 Whole Cell Lysate at 40ug
Predicted bind size: 150KD
Observed bind size: 150KD



Anti- TIF1 gamma Picoband antibody, ABO12522, IHC(P)
IHC(P): Human Placenta Tissue

Anti-TIF1 Gamma Picoband Antibody - Background

Tripartite motif-containing 33 (TRIM33), also known as transcriptional intermediary factor 1 gamma (TIF1- \hat{I}^3), is a human gene. The TRIM33 gene is mapped to chromosome 1p13 by FISH. The protein encoded by this gene is thought to be a transcriptional corepressor. However, molecules that interact with this protein have not yet been identified. The protein is a member of the tripartite motif family. This motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. Three alternatively spliced transcript variants for this gene have been described; however, the full-length nature of one variant has not been determined.