

**Anti-Ceruloplasmin Picoband Antibody**  
Catalog # ABO12538**Specification**

---

**Anti-Ceruloplasmin Picoband Antibody - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">P00450</a>
Host	<b>Rabbit</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Format	<b>Lyophilized</b>

**Description**

Rabbit IgG polyclonal antibody for Ceruloplasmin(CP) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Ceruloplasmin Picoband Antibody - Additional Information**

**Gene ID** 1356

**Other Names**

Ceruloplasmin, 1.16.3.1, Ferroxidase, CP

**Calculated MW**

122205 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat  
<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted. Colocalizes with GCP1 in secretory intracellular compartments. .

**Tissue Specificity**

Expressed by the liver and secreted in plasma.

**Protein Name**

Ceruloplasmin

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

E. coli-derived human Ceruloplasmin recombinant protein (Position: K20-M259). Human Ceruloplasmin shares 80.8% and 79.6% amino acid (aa) sequence identity with mouse and rat Ceruloplasmin, respectively.

**Purification**

Immunogen affinity purified.

#### Cross Reactivity

No cross reactivity with other proteins.

Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

### Anti-Ceruloplasmin Picoband Antibody - Protein Information

Name CP ([HGNC:2295](#))

#### Function

Multifunctional blue, copper-binding (6-7 atoms per molecule) glycoprotein. It has ferroxidase activity oxidizing Fe(2+) to Fe(3+) without releasing radical oxygen species. It is involved in iron transport across the cell membrane (PubMed:[16150804](http://www.uniprot.org/citations/16150804)). Copper ions provide a large number of enzymatic activites. Oxidizes highly toxic ferrous ions to the ferric state for further incorporation onto apo- transferrins, catalyzes Cu(+) oxidation and promotes the oxidation of biogenic amines such as norepinephrin and serotonin (PubMed:[14623105](http://www.uniprot.org/citations/14623105), PubMed:[4643313](http://www.uniprot.org/citations/4643313), PubMed:[5912351](http://www.uniprot.org/citations/5912351)). Provides Cu(2+) ions for the ascorbate-mediated deaminase degradation of the heparan sulfate chains of GPC1 (By similarity). Has glutathione peroxidase-like activity, can remove both hydrogen peroxide and lipid hydroperoxide in the presence of thiols (PubMed:[10481051](http://www.uniprot.org/citations/10481051)). Also shows NO-oxidase and NO2 synthase activities that determine endocrine NO homeostasis (PubMed:[16906150](http://www.uniprot.org/citations/16906150)).

#### Cellular Location

Secreted. Note=Colocalizes with GCP1 in secretory intracellular compartments {ECO:0000250|UniProtKB:P13635}

#### Tissue Location


Expressed by the liver and secreted in plasma.

### Anti-Ceruloplasmin Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Ceruloplasmin Picoband Antibody - Images



Western blot analysis of Ceruloplasmin expression in 22RV1 whole cell lysates (lane 1) and A549 whole cell lysates (lane 2). Ceruloplasmin at 122KD was detected using rabbit anti- Ceruloplasmin Antigen Affinity purified polyclonal antibody at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



Ceruloplasmin was detected in paraffin-embedded sections of human liver cancer tissues using rabbit anti- Ceruloplasmin Antigen Affinity purified polyclonal antibody (Catalog # ABO12538) at 1 µg/mL. The immunohistochemical section was developed using SABC method .

**Anti-Ceruloplasmin Picoband Antibody - Background**

Ceruloplasmin (or caeruloplasmin) is a ferroxidase enzyme that in humans is encoded by the CP gene. It is mapped to 3q23-q25. The protein encoded by this gene is a metalloprotein that binds most of the copper in plasma and is involved in the peroxidation of Fe(II)transferrin to Fe(III) transferrin. Mutations in this gene cause aceruloplasminemia, which results in iron accumulation and tissue damage, and is associated with diabetes and neurologic abnormalities. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene.