

Anti-SPTLC1 Picoband Antibody
Catalog # ABO12580**Specification****Anti-SPTLC1 Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	O15269
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Serine palmitoyltransferase 1(SPTLC1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SPTLC1 Picoband Antibody - Additional Information

Gene ID 10558

Other Names

Serine palmitoyltransferase 1, 2.3.1.50, Long chain base biosynthesis protein 1, LCB 1, Serine-palmitoyl-CoA transferase 1, SPT 1, SPT1, SPTLC1, LCB1

Calculated MW

52744 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Endoplasmic reticulum membrane ; Single-pass membrane protein .

Tissue Specificity

Widely expressed. Not detected in small intestine. .

Protein Name

Serine palmitoyltransferase 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human SPTLC1 (444-473aa IRVVVTVEQTEELERAASTIKEVAQAVLL), different from the related mouse sequence by three amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-SPTLC1 Picoband Antibody - Protein Information

Name SPTLC1

Synonyms LCB1

Function

Component of the serine palmitoyltransferase multisubunit enzyme (SPT) that catalyzes the initial and rate-limiting step in sphingolipid biosynthesis by condensing L-serine and activated acyl-CoA (most commonly palmitoyl-CoA) to form long-chain bases. The SPT complex is also composed of SPTLC2 or SPTLC3 and SPTSSA or SPTSSB. Within this complex, the heterodimer with SPTLC2 or SPTLC3 forms the catalytic core (PubMed: 19416851, PubMed: 33558762, PubMed: 36170811). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed: 19416851, PubMed: 33558762). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1-SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed: 19416851, PubMed: 19648650). The SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC3-SPTSSB isozyme displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed: 19416851, PubMed: 19648650, PubMed: 33558761, PubMed: 33558762). Required for adipocyte cell viability and metabolic homeostasis (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein
{ECO:0000250|UniProtKB:O35704}

Tissue Location


Widely expressed. Not detected in small intestine.

Anti-SPTLC1 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SPTLC1 Picoband Antibody - Images



Western blot analysis of SPTLC1 expression in rat brain extract (lane 1), mouse brain extract (lane 2) and MCF-7 whole cell lysates (lane 3). SPTLC1 at 53KD was detected using rabbit anti-SPTLC1 Antigen Affinity purified polyclonal antibody (Catalog # ABO12580) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



SPTLC1 was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- SPTLC1 Antigen Affinity purified polyclonal antibody (Catalog # ABO12580) at 1 µg/mL. The immunohistochemical section was developed using SABC method .

Anti-SPTLC1 Picoband Antibody - Background

SPTLC1 (Serine palmitoyltransferase, long chain base subunit 1), also known as SPT1, LCB1, is a protein which in humans is encoded by the SPTLC1 gene. Dawkins et al. (2001) noted that the SPTLC1 gene maps to chromosome 9q22.1-q22.3. Serine palmitoyltransferase, which consists of two different subunits, is the initial enzyme in sphingolipid biosynthesis. It converts L-serine and

palmitoyl CoA to 3-oxosphinganine with pyridoxal 5'-phosphate as a cofactor. The product of this gene is the long chain base subunit 1 of serine palmitoyltransferase. Mutations in this gene were identified in patients with hereditary sensory neuropathy type 1. Alternatively spliced variants encoding different isoforms have been identified.