

Anti-CASR Picoband Antibody
Catalog # ABO12609**Specification**

Anti-CASR Picoband Antibody - Product Information

Application	WB, IHC
Primary Accession	P41180
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Extracellular calcium-sensing receptor(CASR) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CASR Picoband Antibody - Additional Information

Gene ID 846

Other Names

Extracellular calcium-sensing receptor, CaSR, Parathyroid cell calcium-sensing receptor 1, PCaR1, CASR, GPRC2A, PCAR1

Calculated MW

120674 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, Rat, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane ; Multi-pass membrane protein .

Tissue Specificity

Expressed in the temporal lobe, frontal lobe, parietal lobe, hippocampus, and cerebellum. Also found in kidney, lung, liver, heart, skeletal muscle, placenta. .

Protein Name

Extracellular calcium-sensing receptor

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E. coli-derived human CASR recombinant protein (Position: Q926-S1078). Human CASR shares 80.5% and 78.6% amino acid (aa) sequence identity with mouse and rat CASR, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r^o Constitution, at 4°C for one month. It Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-CASR Picoband Antibody - Protein InformationName CASR ([HGNC:1514](#))**Function**

G-protein-coupled receptor that senses changes in the extracellular concentration of calcium ions and plays a key role in maintaining calcium homeostasis (PubMed: [17555508](http://www.uniprot.org/citations/17555508), PubMed: [19789209](http://www.uniprot.org/citations/19789209), PubMed: [21566075](http://www.uniprot.org/citations/21566075), PubMed: [22114145](http://www.uniprot.org/citations/22114145), PubMed: [22789683](http://www.uniprot.org/citations/22789683), PubMed: [23966241](http://www.uniprot.org/citations/23966241), PubMed: [25104082](http://www.uniprot.org/citations/25104082), PubMed: [25292184](http://www.uniprot.org/citations/25292184), PubMed: [25766501](http://www.uniprot.org/citations/25766501), PubMed: [26386835](http://www.uniprot.org/citations/26386835), PubMed: [7759551](http://www.uniprot.org/citations/7759551), PubMed: [8636323](http://www.uniprot.org/citations/8636323), PubMed: [8702647](http://www.uniprot.org/citations/8702647), PubMed: [8878438](http://www.uniprot.org/citations/8878438)). Senses fluctuations in the circulating calcium concentration and modulates the production of parathyroid hormone (PTH) in parathyroid glands (By similarity). The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system (PubMed: [7759551](http://www.uniprot.org/citations/7759551)). The G-protein-coupled receptor activity is activated by a co-agonist mechanism: aromatic amino acids, such as Trp or Phe, act concertedly with divalent cations, such as calcium or magnesium, to achieve full receptor activation (PubMed: [27386547](http://www.uniprot.org/citations/27386547), PubMed: [27434672](http://www.uniprot.org/citations/27434672)).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Expressed in the temporal lobe, frontal lobe, parietal lobe, hippocampus, and cerebellum. Also found in kidney, lung, liver, heart, skeletal muscle, placenta.

Anti-CASR Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-CASR Picoband Antibody - Images



Western blot analysis of CASR expression in HELA whole cell lysates (lane 1) and 22RV1 whole cell lysates (lane 2). CASR at 130KD was detected using rabbit anti- CASR Antigen Affinity purified polyclonal antibody (Catalog # ABO12609) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



CASR was detected in paraffin-embedded sections of mouse kidney tissues using rabbit anti-CASR Antigen Affinity purified polyclonal antibody (Catalog # ABO12609) at 1 µg/mL. The immunohistochemical section was developed using SABC method .



CASR was detected in paraffin-embedded sections of rat kidney tissues using rabbit anti- CASR Antigen Affinity purified polyclonal antibody (Catalog # ABO12609) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Anti-CASR Picoband Antibody - Background

The calcium-sensing receptor (CaSR) is a G protein-coupled receptor that is expressed in the parathyroid hormone (PTH)-producing chief cells of the parathyroid gland, and the cells lining the kidney tubule. It senses small changes in circulating calcium concentration and couples this information to intracellular signaling pathways that modify PTH secretion or renal cation handling, thus this protein plays an essential role in maintaining mineral ion homeostasis. Mutations in this gene cause familial hypocalciuric hypercalcemia, familial, isolated hypoparathyroidism, and neonatal severe primary hyperparathyroidism.