

**Anti-GTPase HRAS Antibody**  
Catalog # ABO12775

**Specification**

---

**Anti-GTPase HRAS Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P01112</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for GTPase Hras(HRAS) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-GTPase HRAS Antibody - Additional Information**

**Gene ID** 3265

**Other Names**

GTPase HRas, H-Ras-1, Ha-Ras, Transforming protein p21, c-H-ras, p21ras, GTPase HRas, N-terminally processed, HRAS, HRAS1

**Calculated MW**

21298 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Mouse, Rat, Human<br>

**Subcellular Localization**

Cell membrane. Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus. Golgi apparatus membrane; Lipid-anchor. The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus. .

**Tissue Specificity**

Widely expressed. .

**Protein Name**

GTPase Hras

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human GTPase HRAS

(101-137aa KRVKDSDDVPMVLVGNKCDLAARTVESRQAQDLAR SY), identical to the related mouse and rat sequences.

#### **Purification**

Immunogen affinity purified.

#### **Cross Reactivity**

No cross reactivity with other proteins

#### **Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### **Anti-GTPase HRAS Antibody - Protein Information**

**Name** HRAS

**Synonyms** HRAS1

#### **Function**

Involved in the activation of Ras protein signal transduction (PubMed:[22821884](http://www.uniprot.org/citations/22821884)). Ras proteins bind GDP/GTP and possess intrinsic GTPase activity (PubMed:[12740440](http://www.uniprot.org/citations/12740440), PubMed:[14500341](http://www.uniprot.org/citations/14500341), PubMed:[9020151](http://www.uniprot.org/citations/9020151)).

#### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:P20171}; Lipid-anchor; Cytoplasmic side. Golgi apparatus. Golgi apparatus membrane; Lipid-anchor. Note=The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus.

#### **Tissue Location**


Widely expressed..


### **Anti-GTPase HRAS Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-GTPase HRAS Antibody - Images**





Anti-GTPase HRAS antibody, ABO12775, Western blotting All lanes: Anti GTPase HRAS (ABO12775) at 0.5ug/ml Lane 1: Rat Brain Tissue Lysate at 50ug Lane 2: Mouse Brain Tissue Lysate at 50ug Lane 3: HEPA Whole Cell Lysate at 40ug Predicted bind size: 21KD Observed bind size: 21KD

#### **Anti-GTPase HRAS Antibody - Background**

GTPase HRas, also known as transforming protein p21, is an enzyme that in humans is encoded by the HRAS gene. This gene belongs to the Ras oncogene family, whose members are related to the transforming genes of mammalian sarcoma retroviruses. The products encoded by these genes function in signal transduction pathways. These proteins can bind GTP and GDP, and they have intrinsic GTPase activity. This protein undergoes a continuous cycle of de- and re-palmitoylation, which regulates its rapid exchange between the plasma membrane and the Golgi apparatus. Mutations in this gene cause Costello syndrome, a disease characterized by increased growth at the prenatal stage, growth deficiency at the postnatal stage, predisposition to tumor formation, mental retardation, skin and musculoskeletal abnormalities, distinctive facial appearance and cardiovascular abnormalities. Defects in this gene are implicated in a variety of cancers, including bladder cancer, follicular thyroid cancer, and oral squamous cell carcinoma. Multiple transcript variants, which encode different isoforms, have been identified for this gene.