

Anti-CD40L Picoband Antibody
Catalog # ABO12923

Specification

Anti-CD40L Picoband Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P29965 |
| Host | Rabbit |
| Reactivity | Human |
| Clonality | Polyclonal |
| Format | Lyophilized |

Description

Rabbit IgG polyclonal antibody for CD40L detection. Tested with WB, Direct ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CD40L Picoband Antibody - Additional Information

Gene ID 959

Other Names

CD40 ligand, CD40-L, T-cell antigen Gp39, TNF-related activation protein, TRAP, Tumor necrosis factor ligand superfamily member 5, CD154, CD40 ligand, membrane form, CD40 ligand, soluble form, CD40LG, CD40L, TNFSF5, TRAP

Application Details

Western blot, 0.1-0.5 µg/ml
 Direct ELISA, 0.1-0.5 µg/ml

Subcellular Localization

Cell membrane.

Tissue Specificity

Specifically expressed on activated CD4+ T- lymphocytes.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived human CD40L recombinant protein (Position: E108-L261).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

Anti-CD40L Picoband Antibody - Protein Information

Name CD40LG

Synonyms CD40L, TNFSF5, TRAP

Function

Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:1280226, PubMed:31331973). Costimulates T-cell proliferation and cytokine production (PubMed:8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:8617933). Induces the activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell surface

Tissue Location


Specifically expressed on activated CD4+ T- lymphocytes

Anti-CD40L Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD40L Picoband Antibody - Images






Figure 1. Western blot analysis of CD40L using anti-CD40L antibody (ABO12923).

Anti-CD40L Picoband Antibody - Background

CD40 ligand(CD40L) is a type II membrane protein of 261 amino acids on activated T cells that induces B cell proliferation and immunoglobulin secretion. It has homology with tumour necrosis factor-alpha and -beta, and has important functions in B-cell activation and differentiation. Human CD40L with 5 exons, is mapped to the proximal region of the mouse X chromosome on Xq26.3-27.1, and can be detected on T cells but is absent from B cells and monocytes. Since CD40L is expressed on platelets and released from them on activation, its predictive value as a marker for clinical outcome and the therapeutic effect of inhibition of glycoprotein IIb /IIIa receptor in patients with acute coronary syndromes was investigated. The soluble CD40L may be involved in the process of restenosis and that it exerts its effect by triggering a complex group of inflammatory reactions on endothelial and mononuclear cells. CD40L plays a central role in the pathophysiology of acute coronary syndromes, and has a role in the pathogenesis of coronary artery lesions.